

ON STRING TOPOLOGY OF THREE MANIFOLDS

HOSSEIN ABBASPOUR

ABSTRACT. Let M be a closed, oriented and smooth manifold of dimension d . Let $\mathbb{L}M$ be the space of smooth loops in M . In [5] Chas and Sullivan introduced loop product, a product of degree $-d$ on the homology of $\mathbb{L}M$. In this paper we show how for three manifolds the “nontriviality” of the loop product relates to the “hyperbolicity” of the underlying manifold. This is an application of the existing powerful tool and results in three dimensional topology such as the prime decomposition, torus decomposition, Seifert theorem, torus theorem.

CONTENTS

1. Introduction and the Statement of the Main Theorem	1
2. 3-Manifolds with Finite Fundamental Group	3
3. 3-Manifolds with Non Separating 2-Sphere or 2-Torus	7
4. Closed Seifert Manifolds	9
5. Seifert Manifolds with Boundary	11
6. Connected Sum of 3-Manifolds	15
7. An Injectivity Lemma	16
8. Gluing Seifert Manifolds along Tori	17
9. Gluing to Hyperbolic Manifolds along Torus	22
10. Proof of Main Theorem: Part I	24
11. Proof of Main Theorem: Part II, Algebraically Hyperbolic Manifolds	28
Appendix A. Some Facts in Free Products with Amalgamation	30
Appendix B. Finite Covers of Certain Seifert Manifolds with Boundary	34
References	37

1. INTRODUCTION AND THE STATEMENT OF THE MAIN THEOREM

Throughout this manuscript M denotes a connected, oriented smooth manifold unless otherwise stated. We shall follow the terminology and notations of [5].

We think of the unit circle, $\mathbb{S}^1$, as the quotient $\mathbb{R}/\mathbb{Z}$. The *marked point* of the circle is the image of 0 in this quotient. A *loop* in M is a continuous map from $\mathbb{S}^1$ to M . The *free loop space* $\mathbb{L}M$ of M is the space of all loops in M . If $f : \mathbb{S}^1 \rightarrow M$ is a loop in M then the image of the marked point of $\mathbb{S}^1$ is called the *marked point* of the loop f .

We denote the integral homology of $\mathbb{L}M$ by homology $\mathbb{H}_*(M)$ and is equipped with an *associative product* called *loop product* (see [5]) to be denoted by $\bullet$. The loop product has degree $-d$, where d is the dimension of M namely,

$$\bullet : \mathbb{H}_i(M) \otimes \mathbb{H}_j(M) \rightarrow \mathbb{H}_{i+j-d}(M).$$

Theorem (Chas-Sullivan [5]) For a connected closed oriented smooth manifold M , $(\mathbb{H}_{*-d}(M), \bullet)$ is a graded commutative algebra.

The inclusion of constant loops induces a map

$$i : H_*(M) \rightarrow \mathbb{H}_*(M),$$

and $p : \mathbb{H}_*(M) \rightarrow H_*(M)$ is the map induced by projection on the marked points. Observe that

$$p \circ i = id_{H_*(M)},$$

hence there is a canonical decomposition

$$\begin{aligned} \mathbb{H}_*(M) &= i(H_*(M)) \oplus A_M, \\ A_M &= \text{Ker } p \text{ and } p_{A_M} = id_{\mathbb{H}_*(M)} - p. \end{aligned}$$

Proposition 1.1. (See [5]) $i : H_*(M) \rightarrow \mathbb{H}_*(M)$ is a map of graded algebras where the multiplication on $H_*(M)$ is the intersection product denoted by $\wedge$. Therefore, one can regard $(\mathbb{H}_*(M), \bullet)$ as an extension of $(H_*(M), \wedge)$.

Definition 1.2. We shall say that an oriented closed manifold M has *nontrivial extended loop products* if the restriction of the loop product to A_M is nontrivial, namely

$$\bullet|_{A_M \otimes A_M} \neq 0.$$

Otherwise we shall say that M has *trivial extended loop products*.

Definition 1.3. Let M be a closed 3-manifold. We say that M is *algebraically hyperbolic* if it is a $K(\pi, 1)$ (aspherical) and its fundamental group has no rank 2 abelian subgroup.

Note that any finite cover of an algebraically hyperbolic manifold is also algebraically hyperbolic.

The following is the main theorem of this paper:

Theorem 1.4. *Let M be an oriented closed 3-manifold. If M is not algebraically hyperbolic then M or a double cover of M has nontrivial extended loop products. If M is algebraically hyperbolic then M and all its finite covers have trivial extended loop products.*

From now on we identify $H_*(M)$ with its image under i . The following lemma will be useful to us.

Lemma 1.5. :

- (i) *There is a one-to-one correspondence between the connected components of $\mathbb{L}M$ and the conjugacy classes of $\pi_1(M)$.*
- (ii) *Let $\alpha \in \pi_1(M)$ and let $f \in \mathbb{L}M$ be a loop representing α . Then there exists a short exact sequence*

$$(1.1) \quad \pi_2(M) \cong \pi_1((\Omega M)_\alpha, f) \xrightarrow{S} \pi_1((\mathbb{L}M)_{[\alpha]}, f) \xrightarrow{R} \pi_\alpha \longrightarrow 0$$

where π_α is the centralizer of α in $\pi_1(M)$.

If $[\alpha] \in \hat{\pi}_1(M)$ is a conjugacy class then $(\mathbb{L}M)_{[\alpha]}$ denotes the corresponding connected component of $\mathbb{L}M$ and $\mathbb{H}_*(M)_\alpha$ its integral homology.

If α and β are two loops with the same marked points, their *composition*, $\alpha\beta$, is also a loop and is defined as follows:

$$\alpha\beta = \begin{cases} \alpha(2t) & 0 \leq t \leq 1/2 \\ \beta(2t-1) & 1/2 \leq t \leq 1 \end{cases}$$

We give an informal description of loop product: given two homology classes in $\mathbb{H}_*(M)$, choose a chain representative for each one of them and consider the corresponding set of marked points. They form two chains in M . If these two chains are not in transversal position, replace the representatives with appropriate ones so that their chain of marked points intersect transversally. At each intersection point, compose the corresponding loops of each chain of loops. This process produces a new chain of loops. In [5] it is proved that this operation passes to homology.

Remark 1.6. If M is a closed manifold then the algebra $(\mathbb{H}_*(M), \bullet)$ has a *unit*, μ_M , described as follows. Since each point in M can be regarded as a constant loop, a chain in M gives rise to a chain in $\mathbb{L}M$. This map passes to homology. The image of the fundamental class of M under this map is the unit of $(\mathbb{H}_*(M), \bullet)$.

$\mathbb{H}_*(M)$ is also equipped with a self map of degree 1, denoted Δ , which can be described as follows. Consider the natural action of $\mathbb{S}^1$ on $\mathbb{L}M$: $f \in \mathbb{L}M$ and $r \in \mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$,

$$\begin{aligned} r.f : \mathbb{S}^1 &\rightarrow M \\ (rf)(t) &= f(r+t) \end{aligned}$$

This action induces a map

$$\Delta : \mathbb{H}_*(M) \rightarrow \mathbb{H}_{*+1}(M)$$

Δ has the property that $\Delta^2 = 0$ (see [5]).

Example 1.7. (see [5]) Let M be a closed hyperbolic oriented 3-manifold. As M is aspherical then by Lemma 1.5, each component of $(\mathbb{L}M)_{[\alpha]}$ is a $K(\pi_\alpha, 1)$ where π_α is the centralizer of α in $\pi_1(M)$. If $\alpha \neq 1 \in \pi_1(M)$ then it can be regarded as a hyperbolic element of the discrete subgroup $\pi_1(M)$ of $PSL(2, \mathbb{C})$. So its centralizer is the cyclic group generated by α . Hence $(\mathbb{L}M)_{[\alpha]}$ is a $K(\mathbb{Z}, 1)$ and

$$A_M \simeq H_*\left(\coprod_{[\alpha] \neq [1] \in \hat{\pi}_1(M)} K(\mathbb{Z}, 1)\right).$$

Therefore A_M has homological dimension 1 which implies that the loop product is identically zero on $A_M \otimes A_M$ as a consequence of the absence of homology class of sufficiently large dimension. So closed oriented hyperbolic manifolds have trivial extended loop product.

2. 3-MANIFOLDS WITH FINITE FUNDAMENTAL GROUP

In this section we show that 3-manifolds with finite fundamental group have nontrivial extended loop products.

The following construction is valid for any oriented closed manifold M . Let p be the based point for M . Consider the 0-chain in $\mathbb{L}M$ that consists of the constant loop at p . It represents a homology class $\epsilon \in \mathbb{H}_0(M)$. Given a chain $c \in \mathbb{L}M$ whose set of marked points is transversal to p , then the loops in c whose marked points

are p form a chain in the based loop space of ΩM . This induces a map of degree $-d$

$$\cap : \mathbb{H}_*(M) \rightarrow H_{*-d}(\Omega M).$$

On the other hand $H_*(\Omega M)$ is equipped with Pontryagin product.

Lemma 2.1. (see [5]) $\cap : (\mathbb{H}(M), \bullet) \rightarrow (H_{*-d}(\Omega M), \times)$ is an algebra map where $\times$ is the Pontryagin multiplication. In particular if M is a Lie group then $\cap$ is surjective.

Proposition 2.2. S^3 has nontrivial extended loop products.

Proof. Since S^3 is a Lie group, by Lemma 2.1 $\cap$ is surjective. $(H_*(\Omega S^3), \times)$ is a polynomial algebra with one generator of degree 2. Let $x_1 \in H_p(\Omega S^3)$ and $x_2 \in H_q(\Omega S^3)$ be non-zero elements such that $p, q > 3$. Let $a_i \in \cap^{-1}(x_i)$ for $i = 1, 2$. Since $H_i(S^3) = 0$ for $i > 3$ we have $a_i \in A_{S^3}$. By Lemma 2.1 $\cap(a_1 \bullet a_2) = \cap(a_1) \times \cap_p(a_2) = x_1 \times x_2 \neq 0$, therefore $a_1 \bullet a_2 \neq 0$ $\square$

Lemma 2.3. Let M and N be two closed oriented n -manifolds and $p \in N$. For $f : M \rightarrow N$ which is transversal to p and $f^{-1}(p)$ consists of only one point, the following diagram commutes :

$$(2.1) \quad \begin{array}{ccc} \mathbb{H}_*(M) & \xrightarrow{\cap} & H_*(\Omega M) \\ f_{\mathbb{L}} \downarrow & & f_{\Omega} \downarrow \\ \mathbb{H}_*(N) & \xrightarrow{\cap} & H_*(\Omega N) \end{array}$$

f_{Ω} and $f_{\mathbb{L}}$ are the induced maps by f on the homologies of the based loop space and the free loop space.

Proof. We choose p and $f^{-1}(p)$ respectively as the base point of N and M . We show that the above diagram is commutative even at the chain level. Let $k : \Delta^m \rightarrow \mathbb{L}M$ be a m -simplex in $\mathbb{L}M$. So for each $x \in \Delta^m$

$$k(x) : \mathbb{S}^1 \rightarrow M.$$

If $k_0 : \Delta^m \rightarrow M$,

$$k_0(x) = k(x)(0),$$

is transversal to $f^{-1}(p)$ then consider the set $\Lambda \subset \Delta^m$,

$$\Lambda = \{x \in \Delta^m | k(x)(0) = f^{-1}(p)\}.$$

$k|_{\Lambda}$ is a chain in $\Omega_{f^{-1}(p)}M$ and $\cap(k) = k|_{\Lambda}$. After composing with f , $f \circ k|_{\Lambda}$ is a chain in $\Omega_p N$ and

$$(f_{\Omega} \circ \cap)(k) = f \circ k|_{\Lambda}.$$

On the other hand $f_{\mathbb{L}}(k) = f \circ k$ is a chain in $\mathbb{L}N$. Since k_0 is transversal to $f^{-1}(p)$ and f is transversal to p thus $f \circ k_0$ is transversal to p . Since f^{-1} consists of only one point in M then

$$\{x \in \Delta^m | ((f \circ k)(x))(0) = p\} = \{x \in \Delta^m | k(x)(0) = f^{-1}(p)\} = \Lambda,$$

therefore,

$$(\cap \circ f_{\mathbb{L}})(k) = f \circ k|_{\Lambda} = (f_{\Omega} \circ \cap)(k).$$

$\square$

Lemma 2.4. *Let M be a 3-manifold which is a homotopy 3-sphere and $p_1, p_2, \dots, p_r$ are distinct points in M . Then there is a homotopy equivalence $f : S^3 \rightarrow M$ which is transversal to all p_i 's and $f^{-1}(p_i)$ consists of only one point for each i .*

Proof. Consider a closed ball $D \subset S^3$. Let f be a diffeomorphism from D to a closed ball in $D' \subset M$ where $p_i \in D'$ for all i .

$f|_{\partial D} : \partial D \rightarrow \partial D'$ can be extended to $f : S^3 \setminus \text{Int} D' \rightarrow M \setminus \text{Int} D$ as all the obstructions which are cohomology classes in

$$H^{q+1}(S^3 \setminus \text{Int} D', \partial D', \pi_q(M \setminus \text{Int} D))$$

vanish since $M \setminus \text{Int} D'$ is contractible¹.

Therefore we have a map that $f : S^3 \rightarrow M$ which sends D' to D diffeomorphically and sends the complement of D' to the complement of D and in particular it is transversal to p_i and $f^{-1}(p_i)$ consists of only one point in S^3 . To see that f is a homotopy equivalent it is enough to observe that it has degree one. $\square$

Proposition 2.5. *A closed simply connected 3-manifold M , has nontrivial extended loop products.*

Proof. Let $p \in M$ and $f : S^3 \rightarrow M$ be the homotopy equivalence provided by Lemma 2.4 for $r = 1$.

f induces a homotopy equivalences $f_\Omega : \Omega S^3 \rightarrow \Omega M$ and $f_\mathbb{L} : \mathbb{L} S^3 \rightarrow \mathbb{L} M$. We have the following commutative diagram by Lemma 2.3:

$$(2.2) \quad \begin{array}{ccc} \mathbb{H}_*(S^3) & \xrightarrow{\cap_{S^3}} & H_*(\Omega_{f^{-1}(p)} S^3) \\ f_\mathbb{L} \downarrow & & f_\Omega \downarrow \\ \mathbb{H}_*(M) & \xrightarrow{\cap_M} & H_*(\Omega_p M) \end{array}$$

Since f_Ω and $f_\mathbb{L}$ are isomorphisms and $\cap_{S^3}$ is surjective, therefore $\cap_M$ is also surjective. Let $x_1 \neq 0 \in H_m(\Omega M)$ ² and $x_2 \neq 0 \in H_n(\Omega M)$ where $n, m > 3$ and $a_i \in \cap^{-1}(x_i)$, $i = 1, 2$.

Then, $a_i \in A_M$ since $H_k(M) = 0$ for all $k > 3$. By commutativity of diagram (2.2),

$$f(a_1 \bullet a_2) = f(a_1) \times f(a_2) = x_1 \times x_2 \neq 0.$$

Hence $a_1 \bullet a_2 \neq 0$. $\square$

Proposition 2.6. *If M is a closed oriented 3-manifold with finite fundamental group then M has nontrivial extended loop products.*

Proof. Let $\tilde{M}$ be the universal cover of M and q be the covering map and $r = \text{degree}(q) = |\pi_1(M)| < \infty$. Choose $p \in M$ and let $\{p_1, p_2, \dots, p_r\} \in q^{-1}(p)$ where p_i 's are distinct.

Since $\tilde{M}$ is homotopy equivalent to S^3 , by Lemma 2.4 there is a homotopy equivalence $f : S^3 \rightarrow \tilde{M}$ which is transversal at each p_i and $f^{-1}(p_i)$ consists of only one point in S^3 in M . Let $m_i = f^{-1}(p_i)$, $i = 1, 2, \dots, r$. Hence $(q \circ f)^{-1}(p) = \{m_1, m_2, \dots, m_r\}$ and $q \circ f$ is transversal to p .

¹This can be proved using Mayer-Vietoris exact sequence

² $H_*(\Omega M) \cong H_*(\Omega S^3) \cong \mathbb{Z}[x]$ and degree $x=2$

f composed by the covering map $q : \tilde{M} \rightarrow M$ induces the maps

$$(q \circ f)_{\mathbb{L}} : \mathbb{H}_*(S^3) \rightarrow \mathbb{H}_*(M)$$

and

$$(q \circ f)_{\Omega} : H_*(\Omega_{m_1} S^3) \rightarrow H_*(\Omega_p M)$$

where the latter is an isomorphism.

As we know, generally ΩM and $\mathbb{L}M$ are not connected. So we should emphasize that the images of $(q \circ f)_{\mathbb{L}}$ and $(q \circ f)_{\Omega}$ are homologies of the connected components corresponding to the trivial loop in M and in the diagram below, the sub-index 0 indicates that.

We claim that

$$(2.3) \quad \begin{array}{ccc} \mathbb{H}_*(S^3) & \xrightarrow{r \cdot \cap_{S^3}} & H_*(\Omega_{m_1} S^3) \\ (f \circ q)_{\mathbb{L}} \downarrow & & (f \circ q)_{\Omega} \downarrow \\ (\mathbb{H}_*(M))_0 & \xrightarrow{\cap_M} & H_*((\Omega_p M)_0) \end{array}$$

is commutative. To prove this note that S^3 has a group structure hence

$$\mathbb{L}S^3 \stackrel{homeo.}{\simeq} S^3 \times \Omega S^3,$$

and

$$(2.4) \quad \mathbb{H}_*(S^3) \simeq H_*(S^3) \otimes H_*(\Omega S^3).$$

There are two types of homology classes in $\mathbb{H}_*(S^3)$.

- (1) The classes that under the isomorphism (2.4) correspond to homology classes in $H_0(S^3) \otimes H_*(\Omega S^3)$. These classes can be represented by cycles in $\mathbb{L}S^3$ whose sets of marked points consist of a single point in S^3 .

Type (1) homology classes are mapped to 0 by $\cap_{S^3}$, since we can choose a chain representative whose set of marked point, which consists of one points, is different from all m_i 's. Therefore they are mapped to 0 by $(f \circ q)_{\Omega} \circ (r \cdot \cap_{S^3})$. Indeed their image under $(f \circ q)_{\mathbb{L}}$ have sets of marked point consist of only one point, p . Thus, they are mapped to zero under $\cap_M \circ (f \circ q)_{\mathbb{L}}$. Hence the diagram commutes in this case

- (2) The classes that under isomorphism (2.4) correspond to homology classes in $H_3(S^3) \otimes H_*(\Omega S^3)$. They can be represented by a cycle in $\mathbb{L}S^3$ whose set of marked points is all of S^3 .

Let $\theta = a \otimes b \in \mathbb{H}_3(S^3) \simeq H_3(S^3) \otimes H_*(\Omega S^3)$. Then

$$\cap_{S^3}(\theta) = b$$

so

$$(f \circ q)_{\Omega} \circ (r \cdot \cap_{S^3} \theta) = r(f \circ q)(b)$$

and this is exactly $\cap_M \circ (f \circ q)_{\Omega}(b)$. Note that under $f \circ q$, r copies of b which are based at each one of m_i 's, come together.

Now consider $a_1, a_2 \in \mathbb{H}_*(S^3)$ constructed in the proof of Proposition 2.2. From commutativity of diagram (2.3) and the fact that $(q \circ f)_{\Omega}$ is an isomorphism it follows that

$$(q \circ f)_{\mathbb{L}}(a_1) \bullet (q \circ f)_{\mathbb{L}}(a_2) \neq 0$$

□

3. 3-MANIFOLDS WITH NON SEPARATING 2-SPHERE OR 2-TORUS

Proposition 3.1. $S^1 \times S^2$ has nontrivial extended loop products.

Proof. Consider $a \in S^1$, $b, p \in S^2$ and $p \neq b$ and let (a, p) be the base point of $S^1 \times S^2$. The maps:

$$\alpha : S^1 \rightarrow S^1 \times S^2, \quad \alpha(x) = (x, b),$$

gives rise to a loop whose marked point is (a, b) .

α represents a 0-homology class $\mathbb{H}_0(M)$. The loop $\gamma\alpha\gamma^{-1}$ represents an element $g_1 \in \pi_1(M)$, where γ is a path connecting (a, b) and (a, p) . The free homotopy type of α is $[g_1] \in \hat{\pi}_1(M)$.

Consider the map

$$\beta : S^2 \rightarrow S^1 \times S^2, \quad \beta(y) = (a, y).$$

Also, β can be regarded as an element of $\pi_1(\Omega(S^1 \times S^2)) \simeq \pi_2(S^1 \times S^2)$ or in other words we think of β as a 1-dimensional family of loops. The free homotopy type of the loops of β is the one of the trivial loop. We compose each loop of β with a fixed loop γ' whose marked point is (a, p) and it represents a nontrivial element $g_2 \in \pi_1(S^2 \times S^1)$, $g_2 \neq g_1$. We continue to denote the result by β .




FIGURE 1. A non separating 2-sphere

The image of β under the Hurewicz map

$$\pi_2(\Omega(S^2 \times S^1)) \rightarrow H_2(\Omega(S^2 \times S^1))$$

and then followed by the induced map by inclusion

$$H_2(\Omega(S^2 \times S^1)) \rightarrow \mathbb{H}_2(S^2 \times S^1)$$

gives rise to an element in $\mathbb{H}_2(S^1 \times S^2)$. Again we continue to denote this element by β .

We claim that

$$p_{A_{S^1 \times S^2}}(\Delta\alpha) \bullet p_{A_{S^1 \times S^2}}(\Delta\beta) \neq 0 \in \mathbb{H}_0(S^1 \times S^2)$$

and this proves that $S^1 \times S^2$ has nontrivial extended loop products.

Note that

$$p_{A_{S^1 \times S^2}}(\Delta\alpha) \bullet p_{A_{S^1 \times S^2}}(\Delta\beta) = (\Delta\alpha - p_{S^1 \times S^2}(\Delta\alpha)) \bullet (\Delta\beta - p_{S^1 \times S^2}(\Delta\beta)).$$

In expanding this expression, we get four terms, each one is the loop product of two homology classes, a 1-homology class and a 2-homology class. The set of marked points of the chain representatives of the homology classes involved in

each one of these four terms intersect exactly at one point, namely (a, b) . Hence $p_{A_{S^1 \times S^2}}(\Delta\alpha) \bullet p_{A_{S^1 \times S^2}}(\Delta\beta)$ is equal to the sum of 4 homology classes in $\mathbb{H}_0(S^1 \times S^2)$, each represented by a loop and a $\pm$ sign.

The free homotopy types of these loops are $[g_1 g_2], [g_1], [g_2], 1$ which are different. Therefore the loop product is nontrivial. $\square$

Corollary 3.2. *An oriented closed 3-manifold with a non separating 2-sphere has nontrivial extended loop products.*

Proof. The proof is similar to the one of Proposition 3.1. The only fact we used in the proof of Proposition 3.1 was that there was a non separating 2-sphere (a 2-sphere that meets some closed simple curve only once). Every oriented closed 3-manifold with non separating 2-sphere is indeed a connected sum of $S^2 \times S^1$ with another 3-manifold (see [7]). So $\pi_1(M)$ is the free product of $\pi_1(S^2 \times S^1)$ and the fundamental group of another 3-manifold. Therefore the choices $g_1, g_2 \neq 1, g_2 \neq g_1$ of elements of $\pi_1(M)$, as in the proof of Proposition 3.1, are also possible in this case. $\square$

Proposition 3.3. *If a closed oriented 3-manifold M has a non separating two sided³ incompressible⁴ torus, then M has nontrivial extended loop products.*

Proof. Let $\varphi : S^1 \times S^1 \rightarrow M$ be a non separating torus. H gives rise to a homology class H in $\mathbb{H}_1(M)$, where the marked points are $\varphi(t, 1)$, $t \in S^1$ and the loop passing through $\varphi(t, 1)$ is

$$\varphi_t : S^1 \rightarrow M$$

$$\varphi_t(s) = \varphi(t, s).$$

The loop φ_1 represents an element $h \in \pi_1(M)$ which is nontrivial as the torus is incompressible.




FIGURE 2. A non separating torus

³The normal bundle is trivial.

⁴ π_1 -injective.

Let $\lambda : S^1 \rightarrow M$ be a loop in M which intersects the non separating torus transversally at exactly one point m (see Figure 2). We can assume that the intersection point m is the marked point of λ and we consider this point as the base point of M . Without loss of generality we can assume $m = \varphi(1, 1)$.

λ represents a nontrivial element $l \in \pi_1(M)$ since the homology class it represents is nontrivial.

Also, λ gives rise to a homology class $L \in \mathbb{H}_0(M)$. The free homotopy type of the loops of L is $[l] \in \hat{\pi}_1(M)$.

We claim that

$$p_{A_M}(\Delta L) \bullet p_{A_M}(\Delta H) \neq 0$$

which proves the assertion.

In the expansion of $p_{A_M}(\Delta L) \bullet p_{A_M}(\Delta H) \in \mathbb{H}_0(M)$ we get four terms, each represented by a single loop and $+$ or $-$ sign. The loops have free homotopy type $[l.h]$, $[l]$, $[h]$ and $[1]$. If we show that three out of four terms are different then there cannot be a complete cancellation, thus the product is not trivial.

Assume $[l]$ and $[h]$ are the same. Then the 1-homologies represented by λ and φ_1 are the same. Note that the 1-homology class represented by λ , intersects the 2-homology represented by the torus $\varphi(.,.)$ nontrivially, while the 1-homology class represented by φ_1 intersects the same 2-homology class trivially⁵. This is a contradiction, hence $[l]$ and $[h]$ are different. Similar argument shows that $[l] \neq [1]$. To see that $[h] \neq [1]$ note that the non separating torus does not bound in M . $\square$

4. CLOSED SEIFERT MANIFOLDS

In this section we prove that closed Seifert manifolds have nontrivial extended loop products. Before we state and prove the result we recall the definition of Seifert manifolds.

A 3-manifold is called *Seifert manifold* if it admits a foliation by circles S^1 in such a way that each leaf has a foliated tubular neighborhood $D^2 \times S^1$ which is $D^2 \times S^1$ either with its trivial foliation or its quotient by a standard action of a finite cyclic group $\mathbb{Z}_p$ (acting on D^2 factor). The fibers (leaves) which have a trivially foliated $D^2 \times S^1$ neighborhood are called *normal fibers* and the others are called *singular fibers* and p in above is its *multiplicity*. The space of the fibers forms a surface and it is called the *base surface*.

Notation: Let M be an oriented Seifert manifold with p exceptional fibers and k boundary components. If S , the base surface of M , is orientable, then it has genus $g \in \mathbb{Z}^+$ otherwise S is non orientable and g is the number of cross caps in S . $\pi_1(M)$ has the presentation (see [8]):

(1) If S is orientable and of genus g ,

(4.1)

$$\begin{aligned} \pi_1(M) = \langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_p, d_1, d_2, \dots, d_k, h | & a_i h a_i^{-1} = h, \quad b_i h b_i^{-1} = h, \\ & c_i h c_i^{-1} = h, \quad d_i h d_i^{-1} = h, \\ & c_i^{\alpha_i} h^{\beta_i} = 1 \\ & 1 = \prod_{i=1}^g [a_i, b_i] \prod_{i=1}^p c_i \prod_{i=1}^k d_i \rangle, \end{aligned}$$

⁵Since the torus is two sided (it normal bundle is trivial) and we can push φ_1 off the torus.

(2) If S is nonorientable and has g cross caps.

$$\begin{aligned}
 \pi_1(M) = \langle a_1, \dots, a_g, c_1, \dots, c_p, d_1, d_2, \dots, d_k, h | a_i h a_i^{-1} &= h^{-1}, \\
 c_i h c_i^{-1} &= h^{\delta_i}, \quad d_i h d_i^{-1} = h, \\
 c_i^{\alpha_i} h^{\beta_i} &= 1 \\
 1 &= \prod_{i=1}^g a_i^2 \prod_{i=1}^p c_i \prod_{i=1}^k d_i \rangle
 \end{aligned}
 \tag{4.2}$$

where $0 < \beta_i < \alpha_i$ are integers and $\delta_i = \pm 1$. h corresponds to the normal fiber. Each c_i can be represented by a loop on the base surface going around the singular fiber once. The d_i 's correspond to the boundary components. α_i is the multiplicity of the singular fiber corresponding to c_i and we say that it has type (α_i, β_i) . Note that $\langle h \rangle$ is a normal subgroup of $\pi_1(M)$ and it is central if the S is orientable.

Remark 4.1. An orientable Seifert manifold M may not be oriented as a fibration but it always has a double cover which is Seifert and is orientable as a fibration or in other words its base surface is orientable. This double cover $\tilde{M}$ can be described as following:

$$\tilde{M} = \{(m, o) | m \in M \text{ \& \& } o \text{ an orientation for the fiber passing through } m\}$$

The covering map $\tilde{M} \xrightarrow{q} M$ is

$$q(m, o) = m.$$

The Seifert fibration of $\tilde{M}$ is obtained by pulling back the one of M . If M has boundary the then each boundary component, a torus, gives rise to 2 boundary components each of which is a torus. If M has g cross caps then $\tilde{M}$ has genus $g-1$.

Proposition 4.2. *If M is a closed oriented Seifert 3-manifold then M or a double cover of M has nontrivial extended loop products.*

Proof. By Remark 4.1 we may assume that the base surface of M is orientable. Also we assume $\pi_1(M) \not\cong \mathbb{Z}_2$, since Corollary 2.6 deals with this case. Let h be the generator of $\pi_1(M)$ corresponding to the normal fiber according the presentation (4.1).

There is a natural 3-homology class in $\mathbb{H}_3(M)$ which has a representative whose set of marked points is exactly M and the homotopy type of the loops is $[h]$. The loop associated with a point in M that is on a normal fiber, is the fiber passing through the point. The loop passing through a point on a singular fiber as a map is a multiple of the singular fiber. We denote this homology class by Θ_M . Loops of Θ_M have the free homotopy type $[h]$. We have $p_{A_M}(\Theta_M) = \Theta_M - p_M(\Theta_M) = \Theta_M - \mu_M$ where μ_M is the unit of $\mathbb{H}_*(M)$.

Consider l a loop representing an element $\alpha \in \pi_1(M)$, $\alpha \neq h, 1$. This is possible since we are assuming $\pi_1(M) \neq \mathbb{Z}_2$. Since h is central, the free homotopy class $[\alpha]$ is different from $[h]$ and $[1]$.

α represents a 0-homology class $\hat{\alpha}$ in $\mathbb{H}_0(M)$. We have

$$p_{A_M}(\hat{\alpha}) = \hat{\alpha} - p(\hat{\alpha})$$

where $p(\hat{\alpha})$ is just the constant loop at the marked point of l .

We claim that

$$p_{A_M}(\Theta_M) \bullet p_{A_M}(\hat{\alpha}) \neq 0.$$

In expanding $p_{A_M}(\Theta_M) \bullet p_{A_M}(\hat{\alpha})$ we obtain four terms, all in $\mathbb{H}_0(M)$ and each one is represented by a single loop and a $\pm$ sign. These four loops have free homotopy type $[\alpha.h]$, $[h]$, $[\alpha]$ and $[1]$. Since there are at least three different classes, any linear combination of them with coefficient ± 1 is nonzero. Therefore M has nontrivial extended loop products. $\square$

5. SEIFERT MANIFOLDS WITH BOUNDARY

By torus decomposition, Seifert manifolds with incompressible⁶ boundary are among the building blocks of 3-manifolds. In this section we provide examples of nontrivial extended loop product for various Seifert manifold with boundary. The examples are organized according to the number of boundary components and singular fibers.

Throughout this section M is a compact oriented Seifert manifold with orientable base surface unless otherwise it is stated (see Corollary 5.7).

M has $b \geq 1$ boundary components and p singular fibers and p' denotes the number of the singular fibers of multiplicity greater than 2. We assume ⁷ that

$$b + p \geq 3.$$

g denotes the genus of the base surface of M . In the proofs of the statements in this section g plays no role. So we assume that

$$g = 0.$$

We present two main ideas in constructing nontrivial extended loop products for Seifert manifolds with boundary,

Two curves argument and *Chas' figure eight argument*.

The following table indicates the cases where we apply these arguments.

	argument
$p + b \geq 4$	Two curves argument
$p' + b \geq 3$	Chas' figure eight argument

5.1. $p + b \geq 3$ (Two curves argument).

Proposition 5.1. *Let M be a compact oriented Seifert manifold with $p > 2$ and $b \geq 1$. Then M has nontrivial extended loop products.*

Proof. Let c_1, c_2 and c_3 be the generators of $\pi_1(M)$ corresponding to three singular fibers. Consider a simple curve γ_1 , on the base surface, away from singular fibers and representing the free homotopy class $[c_1 c_2]$. Similarly, consider a simple curve γ_2 on the base surface, away from the singular fibers and representing the free homotopy class $[c_2 c_3]$. Moreover, γ_2 can be chosen such that it has exactly 2 intersection points with γ_1 .

Since γ_1 is away from the singular point, the fibration can be trivialized over γ_1 (see Figure 3). Therefore, we obtain a map

$$f : \mathbb{T}^2 \rightarrow M$$

from the torus to M , where $f(0, \cdot) = \gamma_1$.

⁶ π_1 -injective.

⁷This assumption will be justified later, in section 10, when we recall the torus decomposition.

f gives rise to a homology class $\Gamma_1 \in \mathbb{H}_1(M)$, by declaring $f(t, 0)$, $t \in \mathbb{S}^1$, as the set of marked points. The loop passing through $f(t, 0)$ is $f(t, s)$, $s \in S^1$ and has the free homotopy type $[c_1 c_2]$. Also, the loop γ_2 gives rise to a homology class $\Gamma_2 \in \mathbb{H}_0(M)$.




FIGURE 3. γ_1 and γ_2 and a trivialized fibration over γ_1

We claim that

$$p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M)$$

which proves the proposition.

The sets of marked points of $\Delta\Gamma_1 \in \mathbb{H}_2(M)$ and $\Delta\Gamma_2 \in \mathbb{H}_1(M)$ intersect at two points. Each point contributes four terms in the expansion of $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2)$. All in all eight terms emerge while calculating $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2)$. Each is represented by a conjugacy class in $\hat{\pi}_1(M)$ and a $\pm$ sign. These conjugacy classes are

$$\begin{aligned} & [c_1 c_2 c_3 c_2], [c_1 c_2], [c_2 c_3], [1] \\ & \text{and} \\ & [c_1 c_2^2 c_3], [c_1 c_2], [c_2 c_3], [1]. \end{aligned}$$




FIGURE 4. Two curves argument

Thus in calculating $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \pmod{2}$, only two terms remain, namely (see Figure 4)

$$[c_1 c_2^2 c_3] \quad \text{and} \quad [c_1 c_2 c_3 c_2].$$

To prove the claim it is sufficient to show that these two conjugacy classes are different. For this, consider the group

$$H = \langle c_1, c_2, c_3, d | c_1 c_2 c_3 d_1 = 1, c_i^{\alpha_i} = 1, 1 \leq i \leq 3 \rangle,$$

where d_1 is a generator corresponding to a boundary component. Indeed H is the free product of the groups $\mathbb{Z}_{\alpha_1} * \mathbb{Z}_{\alpha_2} * \mathbb{Z}_{\alpha_3}$.

Consider the homomorphism $\phi : \pi_1(M) \rightarrow H$ which is identity on c_1, c_2, c_3, d_1 and sends the other generators to the trivial element of H . By the presentation (4.1) ϕ is well-defined.

Note that the image of the conjugacy classes $[c_1 c_2^2 c_3]$ and $[c_1 c_2 c_3 c_2]$ under the ϕ are different in $\hat{H}$ as $c_1 c_2^2 c_3$ and $c_1 c_2 c_3 c_2$ are cyclically different reduced words. Therefore $[c_1 c_2^2 c_3]$ and $[c_1 c_2 c_3 c_2]$ are different conjugacy classes in $\pi_1(M)$. $\square$

Remark 5.2. Indeed, in the proof of the proposition above one can replace a singular fiber by a boundary component or in other words a boundary component works as well as a singular fiber of multiplicity zero. To be more explicit, the curves γ_1 and γ_2 can go around a boundary component of the base surface instead of the singular fiber. In the free product H , one of finite cyclic groups is replaced by a infinite cyclic group. The rest of the proof remains the same.

Corollary 5.3. *Let M be a compact oriented Seifert manifold with p singular fibers and q boundary components. If $p + b > 3$ then M has nontrivial extended loop products.*

Proof. If $p \geq 3$ this is just the statement of Proposition 5.1. In the case $p < 3$, by Remark 5.2, in the proof of Proposition 5.1 one replaces the missing singular fiber with one of the extra boundary components. $\square$

5.2. $p' + b \geq 3$, (Chas' figure eight argument). Now we turn to compact oriented Seifert manifolds with the total number of singular fibers and boundary components equals 3.

The idea of the proof of this case is due to Moira Chas. She found this idea while reformulating a conjecture of Turaev (see [4]) on Lie bi-algebras of surfaces, characterizing non self-intersecting closed curves.

Proposition 5.4. *Let M be an oriented Seifert manifold with $p = 2$ singular fibers and $b = 1$ boundary component. Suppose that none of its singular fibers has multiplicity 2. Then M has nontrivial extended loop products.*

Proof. Let c_1 and c_2 be generators of $\pi_1(M)$ corresponding to the singular fibers. Let γ_1 be a smooth curve on the base surface, with one self intersection point, away from singular fibers and representing the free homotopy class $[c_1 c_2^{-1}]$. Similarly, consider a curve γ_2 on the base surface, away from the singular fibers, with one intersection point and representing the free homotopy class $[c_1^{-1} c_2]$. Moreover, γ_2 can be chosen such that it has exactly 2 intersection points with γ_1 .

The fibration can be trivialized over the γ_1 since it is away from the singular point. Therefore, we obtain a map

$$f : \mathbb{T}^2 \rightarrow M$$

from torus to M , where $f(\cdot, 0) = \gamma_1$. This map gives rise to a homology class $\Gamma_1 \in \mathbb{H}_1(M)$, by declaring $f(t, 0)$, $t \in \mathbb{S}^1$, as the set of marked points. The loop

passing through $f(t, 0)$ is $f(t, s)$, $s \in S^1$. All the loops of this family have the free homotopy type $[c_1 c_2^{-1}]$. The loop γ_2 gives rise to a homology class $\Gamma_2 \in \mathbb{H}_0(M)$. We claim that

$$p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M)$$

which proves the proposition.

The set of marked points of $\Delta\Gamma_1 \in \mathbb{H}_2(M)$ and $\Delta\Gamma_2 \in \mathbb{H}_1(M)$ intersect exactly at two points. Each point contributes four terms in the expansion of $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2)$. All in all 8 terms emerge in the expansion of $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2)$. Each one of them is represented by a conjugacy class in $\hat{\pi}_1(M)$ and a $\pm$ sign. These conjugacy classes are

$$\begin{aligned} & [c_1 c_2 c_1^{-1} c_2^{-1}], [c_1^{-1} c_2], [c_1 c_2^{-1}], [1] \\ & \text{and} \\ & [c_1 c_2^{-1} c_1^{-1} c_2], [c_1^{-1} c_2], [c_1 c_2^{-1}], [1]. \end{aligned}$$




FIGURE 5. Figure eight argument

Calculating $p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \pmod 2$, only two terms remain (see Figure 5), namely

$$[c_1 c_2 c_1^{-1} c_2^{-1}] \quad \text{and} \quad [c_1 c_2^{-1} c_1^{-1} c_2].$$

To prove the claim it is sufficient to show that these two homology classes are different. For this, consider the subgroup

$$H = \langle c_1, c_2, d_1 \mid c_1 c_2 d_1 = 1, c_i^{\alpha_i} = 1, i = 1, 2 \rangle$$

where d_1 corresponds to one of the boundary components in presentation (4.1). Indeed H is the free product of cyclic groups generated by c_1 , c_2 and d_1 .

Consider the homomorphism $\phi : \pi_1(M) \rightarrow H$ which is identity on c_1, c_2, d_1 and sends the other generators of $\pi_1(M)$ in presentation (4.1) to the identity element of H . It follows from presentation (4.1) that ϕ is well defined.

Since $\alpha_i \neq 2$, $i = 1, 2$, then $c_i \neq c_i^{-1} \in H$. Thus the images of $c_1 c_2 c_1^{-1} c_2^{-1}$ and $c_1 c_2^{-1} c_1^{-1} c_2$ under ϕ in the free product H are not conjugate as they are cyclically different. Therefore the conjugacy classes $[c_1 c_2 c_1^{-1} c_2^{-1}]$ and $[c_1 c_2^{-1} c_1^{-1} c_2]$ in $\hat{\pi}_1(M)$ are different as well. $\square$

Remark 5.5. The idea of the proof above works unless there is a singular fiber of multiplicity 2 and this case needs a special treatment.

Corollary 5.6. *Let M be a compact oriented Seifert manifold with $p' + b \geq 3$. Then M has nontrivial extended loop products.*

Proof. If $p' \geq 2$ then it is just Proposition 5.4. Otherwise, by Remark 5.2 a boundary component works as well as a singular fiber with multiplicity zero. The proof is the same as of the one of Proposition 5.4 except that one has to replace one of the generators of $\pi_1(M)$ contributed by a singular fiber to a generator corresponding to a boundary component. $\square$

Corollary 5.7. *Let M be a compact oriented Seifert manifold with a non oriented base and $b \geq 2$ boundary components, then M has nontrivial extended loop products.*

Proof. We can modify the proof of Proposition 5.4 considering that by Remark 5.2 a boundary component serves our purposes as well as a singular fiber. Note that the fibration is oriented once restricted to the boundary components. So it can be trivialized over a simple curve representing the free homotopy type $[d_1 d_2^{-1}]$ where d_1 and d_2 are generators of $\pi_1(M)$ contributed by the two boundary components (see the presentation 4.2). The rest of the proof is similar to the one of Proposition 5.4. Calculating mod 2 we get two terms

$$[d_1 d_2 d_1^{-1} d_2^{-1}] \quad \& \quad [d_1 d_2^{-1} d_1^{-1} d_2].$$

If a_1 is a generators of $\pi_1(M)$ contributed by a cross cap (see the presentation 4.2) then consider the group

$$H = \langle d_1, d_2, a_1 | d_1 d_2 a_1^2 = 1 \rangle \simeq \langle d_1 \rangle * \langle a_1 \rangle$$

and the homomorphism $\varphi : \pi_1(M) \rightarrow H$ which is identity on d_1, d_2, a_1 and sends the other generators of $\pi_1(M)$. The image of the two conjugacy classes above in the free product H is $[d_1 a_1^2 d_1^{-1} a_1^{-2}] \quad \& \quad [d_1 a_1^{-2} d_1^{-1} a_1^2]$ which are different. Therefore the conjugacy classes above are distinct conjugacy classes of $\pi_1(M)$. $\square$

6. CONNECTED SUM OF 3-MANIFOLDS

In this section we consider the connected sum of 3-manifolds. We shall show how one can obtain nontrivial loop products in this kind of manifold. For this part, the author has benefited from conversations with numerous colleagues [1].

Proposition 6.1. *If M_1 and M_2 are two 3-manifolds such that $\pi_1(M_i) \neq 1$, $i = 1, 2$, then $M_1 \# M_2$ has nontrivial extended loop products.*

Proof. Let $m_i \in M_i$, $i = 1, 2$ be two points which are identified in the connected sum $M_1 \# M_2$ and considered as the base point m of M_1 , M_2 and $M_1 \# M_2$. Let $g_1 \in \pi_1(M_1)$ and $g_2 \in \pi_1(M_2)$ be two non-trivial elements. Consider the loops γ_i in M_i with marked point m and representing g_i , $i = 1, 2$.

By composing these two loops at m , we obtain a loop $\gamma_1 \gamma_2$ representing $g_1 g_2 \in \pi_1(M_1 \# M_2) \simeq \pi_1(M_1) * \pi_1(M_2)$ and it has the free homotopy type $[g_1 g_2] \in \hat{\pi}_1(M)$ (see Figure 6). Notice that $\gamma_1 \gamma_2$ also represents an element of $\mathbb{H}_0(M_1 \# M_2)$. We denote this homology class by $g_1 \# g_2$.

Consider the embedding $\beta : S^2 \hookrightarrow M_1 \# M_2$ where the connected sum happens (see Figure 6). This gives rise to an element in $\pi_2(M_1 \# M_2) \simeq \pi_1(\Omega(M_1 \# M_2))$ and therefore an element in $\pi_1(\mathbb{L}(M_1 \# M_2))$. The image of this element under the Hurewicz map is an element of $\mathbb{H}_1(M_1 \# M_2)$. By composing the loops of this 1-cycle of loops with a fixed loop whose marked point is m and represents a nontrivial element h in $\pi_1(M_1 \# M_2)$, we can obtain a new element in $\mathbb{H}_1(M_1 \# M_2)$. We denote this element by $\beta_{M_1 \# M_2}$ and the homotopy type of its loops is $[h] \in \hat{\pi}_1(M_1 \# M_2)$.




FIGURE 6. Loop product in connected sum

We claim that

$$p_{A_M}(\Delta(g_1 \# g_2)) \bullet p_{A_M}(\Delta(\beta_{M_1 \# M_2})) \neq 0$$

which proves that M has nontrivial extended loop product. To compute

$$(6.1) \quad p_{A_M}(\Delta(g_1 \# g_2)) \bullet p_{A_M}(\Delta(\beta_{M_1 \# M_2})) = (\Delta(g_1 \# g_2) - p_M(\Delta(g_1 \# g_2))) \bullet (\Delta(\beta_{M_1 \# M_2}) - p_M(\Delta(\beta_{M_1 \# M_2})))$$

one has to homotop $\gamma_1 \gamma_2$ to a simple curve γ which intersects the embedded S^2 in $M_1 \# M_2$ at two different points. So in calculating the expansion above we get eight terms, four terms for each intersection point of γ and the embedded 2-sphere. Each one of these eight terms, which are elements of $\mathbb{H}_0(M)$, is represented by a single loop and a $\pm$ sign. These loops have free homotopy types,

$$[g_1 g_2 h], [g_1 g_2], [h], [1], [g_2 g_1 h], [g_2 g_1], [h], [1].$$

So calculating $p_{A_M}(\Delta(g_1 \# g_2)) \bullet p_{A_M}(\Delta(\beta_{M_1 \# M_2})) \bmod 2$, only two terms remain,

$$[g_1 g_2 h] \& [g_2 g_1 h].$$

If we choose $h = g_1 g_2$ in $\pi_1(M_1 \# M_2)$, the two conjugacy classes

$$[g_1 g_2 h] = [g_1 g_2 g_1 g_2] \text{ and } [g_2 g_1 h] = [g_1 g_2^2 g_1] = [g_2^2 g_1^2]$$

are different as two cyclically reduced sequences (g_1, g_2, g_1, g_2) and (g_2^2, g_1^2) are not cyclic permutation of one another.

This proves that for $h = g_1 g_2$, $p_{A_M}(\Delta(g_1 \# g_2)) \bullet p_{A_M}(\Delta(\beta_{M_1 \# M_2}))$ is nonzero mod 2, therefore it is nonzero in $\mathbb{H}_0(M_1 \# M_2)$. $\square$

7. AN INJECTIVITY LEMMA

Suppose that M is a closed oriented manifold and $\mathcal{T}$ is a collection of incompressible⁸ separating tori in M . Let $M \setminus \mathcal{T} = M_1 \cup M_2 \dots \cup M_n$. We associate with $(M, \mathcal{T})$ a tree of groups (G, T) (See Appendix A for the definitions).

The vertices of T are in one-to-one correspondence with M_i 's. Two vertices v_i and v_j are connected by an edge if the corresponding M_i and M_j are glued along a torus. With each vertex v_i we associate the fundamental group of $\pi_1(M_i)$ and with each edge we associate the fundamental group of the corresponding torus. As the tori are incompressible there are two monomorphisms from each edge group to the vertex groups of the corresponding vertices. Namely if e is an edge (a torus in M) and v a vertex of e (some M_i s.t. $T \subset \partial \overline{M_i}$) then there is an homomorphism

$$f_e^v : G_e \rightarrow G_v$$

⁸ π_1 -injective

which is indeed the map induced by inclusion

$$i : \pi_1(T) \rightarrow \pi_1(\overline{M_i}) \simeq \pi_1(M_i).$$

By Van-Kampen theorem, $\pi_1(M)$ is the amalgamation of G_v 's along G_e 's (see Appendix II, tree of groups).

Combining all these with Lemma A.5 in Appendix A, we have,

Lemma 7.1. (*Injectivity Lemma*) *Let M , $\mathcal{T}$ and M_i 's be as above. Suppose that $[a]$ and $[b]$ are two distinct conjugacy classes in $\pi_1(M_i)$ such that a is not conjugate to the image of any element of the edge groups in $\pi_1(M_i)$. Then a and b represent distinct conjugacy classes of $\pi_1(M)$.*

The injectivity lemma will be very useful to show that certain examples of non-trivial loop products in a submanifold of M give rise to nontrivial loop product in M .

8. GLUING SEIFERT MANIFOLDS ALONG TORI

In this section we consider the gluing of the Seifert manifolds along tori. We prove that the result of gluing Seifert manifolds along incompressible *separating* tori has nontrivial extended loop products except certain classes for which there are double covers that have nontrivial extended loop products.

The idea of the proof is to apply either the *two curves argument* or the *figure eight argument* to a Seifert piece of the gluing that fulfills the condition of the arguments. Then we show that the examples of nontrivial loop product in the piece give rise to a nontrivial example of loop product in the manifold, for that we need the injectivity lemma, Lemma 7.1.

When we cannot apply either the two curves argument or the figure eight argument we consider an appropriate double cover of the manifold which either has a non separating torus or one of the two arguments can be applied to one of the pieces in its torus decomposition⁹.

Before stating the result and demonstrating the proof we introduce the following notation.

Convention and Notation: Let M be a closed 3-manifold and $\mathcal{T} \neq \emptyset$ be a collection of tori in M and

$$M \setminus \mathcal{T} = M_1 \cup M_2 \cdots \cup M_n,$$

where all the $\overline{M_i}$'s are oriented Seifert manifolds with incompressible boundary¹⁰.

Let $b_i \geq 1$ and p_i denote respectively the number of boundary components and singular fibers of $\overline{M_i}$ and $A_i \subset \mathbb{Z}^+$ is the set of all the multiplicities of the singular fibers of $\overline{M_i}$. The base surface of M_i is denoted S_i and the genus of S_i is g_i if S_i is oriented otherwise g_i is the number of cross caps in S_i .

We assume that all tori in T are separating since in Section 3 we proved that 3-manifolds with non separating torus have nontrivial extended loop products.

We assume¹¹ that for all i

$$b_i + p_i \geq 3$$

⁹See Section 10 for the definition of torus decomposition

¹⁰We say that the torus decomposition of M has only Seifert pieces.

¹¹see section 10 for justification.

if S_i is orientable. Also we assume that

$$g_i = 0$$

if S_i is orientable, as if $g_i > 0$ then there is a non separating torus in M and this case has already been studied in Section 3.

Proposition 8.1. *Let M be a closed 3-manifold described as above. Then M or a double cover of M has nontrivial extended loop products.*

Proof. M belongs to one of the following categories:

- (1) $b_i \geq 3$ and S_i is orientable for some i . In this case we use the figure eight argument. Note that if $b_i > 3$ we can use the two curves argument.
- (2) $b_i \geq 2$ and S_i is not orientable for some i , for this case we also use the figure eight argument .
- (3) $p_i + b_i > 3$ and S_i orientable for some i , we can apply two curves argument.
- (4) $b_i + p_i = 3$, $1 \leq b_i \leq 2$, S_i orientable and $2 \notin A_i$ for some i . In this case we use figure eight argument.
- (5) For all i , $b_i + p_i = 3$, $1 \leq b_i \leq 2$, S_i orientable and $2 \in A_i$ for all i . For this case we consider certain double cover of M .
- (6) $b_i + p_i = 3$ and $2 \in A_i$ if S_i is orientable otherwise $b_i = 1$ and S_i is not orientable. In this case also we consider a double cover.

We verify the statement for each cases.

- (1) **$b_i \geq 3$ and S_i orientable for some i :**

Suppose that $b_1 \geq 3$. By Corollary 5.6 $\overline{M_1}$ has nontrivial extended loop products. Consider the homology classes Γ_1 and $\Gamma_2 \in \mathbb{H}_1(\overline{M_1})$ constructed in the proof of Corollary 5.6. Γ_1 and Γ_2 can be regarded as the elements of $\mathbb{H}_*(M)$. We claim that

$$p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M).$$

In calculating the product above, four terms emerge, all of them in $\mathbb{H}_0(M)$. To prove the product is nonzero it is sufficient to show it is nonzero mod 2. Having done the calculation mod 2, just like in the proof of Proposition 5.4 and Corollary 5.6, only two terms survive,

$$[d_1 d_2 d_1^{-1} d_2^{-1}] \text{ and } [d_1 d_2^{-1} d_1^{-1} d_2],$$

where d_1 and d_2 are the generators contributed by two boundary components (see presentation (4.1)).

By the proof of Corollary 5.6 (indeed Proposition 5.4) we know these two are distinct conjugacy classes of $\pi_1(M_1)$. So if we show that $d_1 d_2 d_1^{-1} d_2^{-1}$ is not conjugate to any element of the fundamental group of one of the components of $\partial\overline{M_1}$, then by injectivity lemma $[d_1 d_2 d_1^{-1} d_2^{-1}]$ and $[d_1 d_2^{-1} d_1^{-1} d_2]$ are different as the conjugacy classes of $\pi_1(M)$.

Suppose that

$$(8.1) \quad [d_1 d_2 d_1^{-1} d_2^{-1}] = [h^r d^s]$$

where h is the generator of $\pi_1(M_1)$ that corresponds to a normal fiber of M_1 and d is a generator of $\pi_1(M)$ contributed by a boundary component.

Consider the group H

$$H = \langle d_1, d_2, d_3 | d_1 d_2 d_3 = 1 \rangle \simeq \langle d_1 \rangle * \langle d_2 \rangle$$

and the homomorphism

$$\phi : \pi_1(M_1) \rightarrow H$$

which is identity on d_1, d_2, d_3 and trivial on all other generators of $\pi_1(M_1)$. It follows from the presentation (4.1) of $\pi_1(M_1)$ that ϕ is well-defined.

After applying ϕ to (8.1) we get

$$[d_1 d_2 d_1^{-1} d_2^{-1}] = [1]$$

if $d \neq d_1, d_2, d_3$ otherwise

$$[d_1 d_2 d_1^{-1} d_2^{-1}] = [d^s].$$

The first case is impossible as $d_1 d_2 d_1^{-1} d_2^{-1}$ is a cyclically reduced word in the free product $\langle d_1 \rangle * \langle d_2 \rangle$ thus it represents a nontrivial conjugacy class.

In the latter case, If $d = d_1$ (or d_2) then

$$[d_1 d_2 d_1^{-1} d_2^{-1}] = [d_1^s],$$

and this is not possible either as $d_1 d_2 d_1^{-1} d_2^{-1}$ is cyclically reduced of length 4 and d_1^s (or d_2^s) has length 1, hence they represent different conjugacy classes.

If $d = d_3 = d_2^{-1} d^{-1}$ then we have

$$[d_1 d_2 d_1^{-1} d_2^{-1}] = [(d_1 d_2)^{-s}].$$

Considering the length of the word we have to have $s = \pm 2$. In either case this inequality does not hold as $d_1 \neq d_1^{-1}$ and $d_2 \neq d_2^{-1}$ (H is a free group). Therefore these two conjugacy classes are always distinct. This proves the claim.

(2) $b_i \geq 2$ and S_i is not orientable:

Similar to the previous case we use the figure eight argument in M_i . Suppose that $b_1 \geq 2$ and S_1 is not orientable. By Corollary 5.7 $\overline{M_1}$ has non-trivial extended loop products. The homology classes Γ_1 and Γ_2 in $\mathbb{H}_*(\overline{M_1})$ constructed in the proof of Corollary 5.7 can be regarded as homology classes of $\mathbb{H}_*(\overline{M})$. We claim that

$$p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M).$$

Calculating mod 2 we get two terms represented by the conjugacy classes $[d_1 a_1^2 d_1^{-1} a_1^{-2}]$ and $[d_1 a_1^{-2} d_1^{-1} a_1^2]$ (see the proof of Corollary 5.7). We must prove that these two conjugacy are distinct as conjugacy classes of M . For that, by injectivity lemma, it is sufficient to prove that one of them has no boundary representative. We show that $[d_1 a_1^2 d_1^{-1} a_1^{-2}]$ has no boundary representative. Suppose we have

$$(8.2) \quad [d_1 a_1^2 d_1^{-1} a_1^{-2}] = [h^r d^s].$$

where d is a generator of $\pi_1(M_1)$ contributed by a boundary component and h corresponds to the normal fiber. Consider the group

$$H = \langle d_1, d_2, a_1 | d_1 d_2 a_1^2 = 1 \rangle \simeq \langle d_1 \rangle * \langle a_1 \rangle \simeq \langle d_2 \rangle * \langle a_1 \rangle$$

and the homomorphism $\varphi : \pi_1(M_1) \rightarrow H$ which is identity on d_1, d_2, a_1 and trivial on the other generators of $\pi_1(M_1)$. It follows from the presentation (4.2) that φ is well defined. Applying φ to (8.2) we have

$$[d_1 a_1^2 d_1^{-1} a_1^{-2}] = [1]$$

if $d \neq d_1, d_2$, otherwise

$$[d_1 a_1^2 d_1^{-1} a_1^{-2}] = [d^s]$$

in the free product $H \simeq \langle d_1 \rangle * \langle a_1 \rangle$. The first case is impossible as $d_1 a_1^2 d_1^{-1} a_1^{-2}$ is a cyclically reduced word of length 4 so it represents a nontrivial conjugacy class. If $d = d_1$ then

$$[d_1 a_1^2 d_1^{-1} a_1^{-2}] = [d_1^s]$$

which is again impossible in the free product $\langle d_1 \rangle * \langle a_1 \rangle$ as the cyclically reduced words $d_1 a_1^2 d_1^{-1} a_1^{-2}$ and d_1^s are different in the free product $\langle d_1 \rangle * \langle a_1 \rangle$. If $d = d_2$ then

$$[d_1 a_1^2 d_1^{-1} a_1^{-2}] = [(a_1^2 d_1)^{-s}].$$

Again similar to previous cases we must have $s = \pm 2$. Even this is not enough as $d_1 \neq d_1^{-1}$ and $a_1^2 \neq a_1^{-2}$.

(3) $p_i + b_i \geq 4$ and S_i **orientable for some i** :

Since case 1 deals with the case $b_i \geq 3$ for some i so we may assume that $b_i \leq 2$. So we are dealing with one of the following cases:

- (i) $p_i > 2$ for some i
- (ii) $b_i = p_i = 2$ for some i .

The proof of both cases are similar and use the two curves argument. We just have to remember that a boundary component works as a singular fiber of multiplicity 0. Here we only present the proof of case (i).

Suppose that $p_1 > 2$. Consider the homology classes Γ_1 and Γ_2 in $\mathbb{H}_*(\overline{M}_1)$ introduced in the proof of Proposition 5.1. They can be considered as the elements of $\mathbb{H}_*(M)$. We claim that

$$p_{A_M}(\Delta\Gamma_1) \bullet p_{A_M}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M)$$

The calculation is the same and the results is a sum of conjugacy classes of $\pi_1(M)$. Computing mod 2 only two terms

$$[c_1 c_2 c_3 c_2] \text{ and } [c_1 c_2^2 c_3].$$

We must prove that they are different conjugacy classes of $\pi_1(M)$. By the proof of Proposition 5.1 we know that they are distinct as the conjugacy classes of $\pi_1(M_1)$. If we show that at least one of them has no representative in the fundamental group of any of the boundary components of M_1 then by injectivity lemma they are different as conjugacy classes of $\pi_1(M)$. Suppose that

$$(8.3) \quad [c_1 c_2 c_3 c_2] = [h^r d^s]$$

where h stands for the normal fiber and d is the generator coming from a boundary component.

Consider the group,

$$H = \langle c_1, c_2, c_3 \mid c_1 c_2 c_3 = 1, c^{\alpha_i} = 1, 1 \leq i \leq 3 \rangle,$$

where α_i is the multiplicity of the singular fiber corresponding to c_i , and the homomorphism $\phi : \pi_1(M_1) \rightarrow H$ which is identity on c_1, c_2, c_3 and sends the other generators of $\pi_1(M_1)$ to the trivial element of H . Applying ϕ to (8.3) we have

$$[c_1 c_2 c_3 c_2] = [1]$$

or

$$c_1 c_2 c_3 c_2 = 1$$

in H . Since $c_1 c_2 c_3 = 1$ in H we conclude that $c_2 = 1$ which is a contradiction.

- (4) $b_i + p_i = 3$, $1 \leq b_i \leq 2$, S_i **orientable** and $2 \notin A_i$ **for some i** :

Here we shall use the figure eight argument. Suppose $i = 1$. By assumption, one of the following holds:

$$(b_1 = 1 \text{ and } p_1 = 2) \text{ or } (b_1 = 2 \text{ and } p_1 = 1.)$$

The proof is similar for both cases, considering that a boundary component behaves just like a singular fiber of multiplicity 0. So we only present the proof of the first case.

Suppose $p_1 = 2$ and $b_1 = 1$. Consider the homology classes Γ_1 and $\Gamma_2 \in \mathbb{H}_*(\overline{M_1})$ as constructed in the proof of the Proposition 5.4. They can be regarded as homology classes in $\mathbb{H}_*(M)$. We prove that

$$p_{AM}(\Delta\Gamma_1) \bullet p_{AM}(\Delta\Gamma_2) \neq 0 \in \mathbb{H}_0(M)$$

which proves that M has nontrivial extended products.

Having done the same mod 2 calculation as the proof of Proposition 5.4 we get two conjugacy classes

$$[c_1 c_2 c_1^{-1} c_2^{-1}] \quad \text{and} \quad [c_1 c_2^{-1} c_1^{-1} c_2].$$

Just like the previous cases, in order to show that these two conjugacy classes are distinct, we must prove that at least one of them has no representative in the fundamental group of the boundary component.

Suppose that

$$(8.4) \quad [c_1 c_2 c_1^{-1} c_2^{-1}] = [h^r d^s]$$

where h is the generator corresponding to the normal fiber and d is the generator corresponding to a boundary component.

Consider the group

$$H = \langle c_1, c_2, d_1 \mid c_1 c_2 d_1 = 1, c_1^{\alpha_1} = c_2^{\alpha_2} = 1 \rangle$$

and the homomorphism $\phi : \pi_1(M_1) \rightarrow H$ which is identity on c_1, c_2, d_1 and sends other generators of $\pi_1(M)$ to the trivial element of H . It follows from presentation (4.1) that ϕ is well-defined.

By applying ϕ to (8.4) we have

$$[c_1 c_2 c_1^{-1} c_2^{-1}] = [d_1^s].$$

in H , or

$$[c_1 c_2 c_1^{-1} c_2^{-1}] = [(c_1 c_2)^s].$$

Note that H is the free product $\mathbb{Z}_{\alpha_1} * \mathbb{Z}_{\alpha_2}$ with c_1 and c_2 as the generators of the corresponding factors.

So if $c_1 c_2 c_1^{-1} c_2^{-1}$ and $(c_1 c_2)^s$ are conjugate then $s = 2$. Even if $s = 2$ then two cyclically reduced words $c_1 c_2 c_1^{-1} c_2^{-1}$ and $c_1 c_2 c_1 c_2$ are cyclically different since c_1 and c_2 are not of multiplicity 2.

- (5) $b_i + p_i = 3$, $1 \leq b_i \leq 2$, S_i **orientable** and $2 \in A_i$ **for all i** :

This is a case where the figure eight argument does not work. The following lemma provides us a double cover of the manifold which has nontrivial extended products by previous cases or the non separating torus argument. Appendix B is devoted to the proof of this lemma.

Lemma 8.2. *Let M be as above, $|\mathcal{T}| = n$, $b_i + p_i = 3$, $1 \leq b_i \leq 2$ and $2 \in A_i$ and S_i orientable for all i .*

- (i) *If $n \geq 3$ then M has a double cover with no non separating torus whose torus decomposition has a Seifert piece with 3 boundary components.*
 - (ii) *If $n = 2$ and $r \in A_1$, $r \neq 2$, then M has a double cover with no non separating torus whose torus decomposition has a Seifert piece with two singular fibers of multiplicity $r \neq 2$.*
 - (iii) *If $n = 2$ and $A_1 = A_2 = \{2\}$ then M has a double cover with a non separating torus.*
- (6) $n = |\mathcal{T}|$:
- $b_i = 2$, $p_i = 1$, $A_i = \{2\}$ **and S_i is orientable for $i \neq 1, n$**
 - $b_1 = b_n = 1$;
 - S_1 **is non orientable**;
 - **If S_n is orientable then $p_n = 1$ and $2 \in A_n$**

This case is similar to the previous case except that some $\overline{M}_i$ may not be orientable as fibration.

If $\overline{M}_1$ has a singular fiber or S_1 has more than one cross cap: then consider M'_1 the double cover of $\overline{M}_1$ which is Seifert manifold and is oriented as fibration (see Remark 4.1). M'_1 has at least 2 singular fibers and 2 boundary components or its base surface has genus greater than zero. We can construct $\tilde{M}$, a double cover of M , by gluing two copies of $M \setminus M_1$ to M'_1 along the boundary components of M'_1 such that the result double covers M . The covering on the complement of M'_1 is the trivial double cover. Note that the torus decomposition of $\tilde{M}$ has a Seifert piece M'_1 which has either two boundary components and 2 singular fibers thus by case (3) $\tilde{M}$ has nontrivial extended loop products or its base surface has genus greater than 0 thus it has a non separating torus and therefore nontrivial extended loop products.

If $\overline{M}_1$ has no singular fiber and S_1 has one cross cap: then S_1 is indeed Möbius strip and $\overline{M}_1$ has another Seifert fibration model whose base surface is a disk and has 2 singular fibers of multiplicity 2. Now if S_n is orientable this is just case (5). If S_n is not orientable then it is the case above.

□

9. GLUING TO HYPERBOLIC MANIFOLDS ALONG TORUS

In this section we analyze gluing of Seifert/hyperbolic and hyperbolic 3-manifolds along torus and for that we need the following lemma. The proof can be found in Appendix A (see A.4).

Lemma 9.1. Suppose that G_1, G_2 and H are three groups with monomorphisms $\phi_i : H \rightarrow G_i$, $i = 1, 2$ and $G_1 *_H G_2$ denotes the amalgamated free product of G_1 and G_2 over H . Let $g_1 \in G_1 \setminus H$ and $g_2 \in G_2 \setminus H$ and $h \neq 1$ in H such that:

- (i) $g_1^{-1} H g_1 \cap H = 1$
- (ii) $g_2 h \neq h g_2$.

Then two conjugacy classes $[h g_1 g_2]$ and $[h g_2 g_1]$ of $G_1 *_H G_2$ are different.

Proposition 9.2. Let M be a connected oriented closed 3-manifold which contains an incompressible torus T such that:

- (i) $M - T$ is disconnected with two components M_1 and M_2 .
- (ii) $\overline{M_1}$ has a hyperbolic interior with finite volume.
- (iii) $\overline{M_2}$ either has a hyperbolic interior with finite volume or is Seifert fibred not homeomorphic to $S^1 \times S^1 \times I$.

Then M has nontrivial extended loop products.

Proof. Let $\varphi : S^1 \times S^1 \rightarrow T \subset M$ be a homeomorphism. φ gives rise to an element $H \in \mathbb{H}_1(M)$, whose marked points are $\varphi(s, 1)$, $s \in S^1$ and the loop passing through $\varphi(s, 1)$ is

$$\begin{aligned} \varphi_s : S^1 &\rightarrow M \\ \varphi_s(t) &= \varphi(s, t) \end{aligned}$$

Take $\varphi(1, 1)$ as the base point for M , $\overline{M_1}$ and $\overline{M_2}$. Let $h = [\varphi_1] \in \pi_1(M)$.




FIGURE 7. M_1 and M_2 glued along T

We recall that $\pi_1(M)$ is the amalgamated free product of $\pi_1(\overline{M_1})$ and $\pi_1(\overline{M_2})$ over $\pi_1(T)$. Pick two arbitrary elements $g_1 \in \pi_1(\overline{M_1})$ and $g_2 \in \pi_1(\overline{M_2})$. Let $g_1 g_2 \in \pi_1(M)$ be their product in the amalgamated free product $\pi_1(\overline{M_1}) *_\pi_1(T) \pi_1(\overline{M_2})$. Let γ_1 and γ_2 be two loops in $\overline{M_1}$ and $\overline{M_2}$ representing g_1 and g_2 . The composition $\gamma_1 \gamma_2$ of these two loops is a representative for $g_1 g_2$. This loop has free homotopy type $[g_1 g_2]$ (see Figure 7).

Replace $\gamma_1 \gamma_2$ by a loop γ with the same free homotopy type which intersects transversally the torus T at two points (see Figure 7). γ can be regarded as an element of $\mathbb{H}_0(M)$. We denote this homology class by Γ .

We claim that there is a choice of $g_i \in \pi_1(\overline{M_i}) \setminus \pi_1(T)$, $i = 1, 2$, and φ such that:

$$p_{A_M}(\Delta \Gamma) \bullet p_{A_M}(\Delta H) \neq 0.$$

In the expansion of $p_{A_M}(\Delta \Gamma) \bullet p_{A_M}(\Delta H)$, we get 4 terms for each intersection point, just like in the proof of Proposition 3.3, therefore 8 terms overall. Each term

is an element of $\mathbb{H}_0(M)$ and represented by a loop. They have the free homotopy types:

$$[hg_1g_2], [h], [g_1g_2], 1, [hg_2g_1], [h], [g_1g_2], 1.$$

Calculating mod 2, we get $[hg_1g_2] + [hg_2g_1]$. By Lemma 9.1 $[hg_1g_2]$ and $[hg_2g_1]$ are different conjugacy classes if $\pi_1(\overline{M_1})$, $\pi_1(\overline{M_2})$ and $\pi_1(T)$ satisfy the conditions of Lemma 9.1.

By assumption the interior of $\overline{M_1}$ is hyperbolic. $\pi_1(\overline{M_1})$, can be regarded as a subgroup of $PSL(2, \mathbb{C})$ where the image of $\pi_1(T)$ in $\pi_1(\overline{M_1})$ consists of parabolic elements which have a common fixed point.

For $g \in \pi_1(\overline{M_1}) \setminus \pi_1(T)$ (we have denoted the image of $\pi_1(T)$ in $\pi_1(\overline{M_1})$ simply by $\pi_1(T)$), $g^{-1}\pi_1(T)g \cap \pi_1(T) = \{1\}$, since if the fixed point of the elements of $\pi_1(T)$ is m then the elements of $g^{-1}\pi_1(T)g$ fix $g^{-1}(m)$. So if $g^{-1}\pi_1(T)g \cap \pi_1(T)$ is nontrivial then m must be a fixed point of g but this implies that $g \in \pi_1(T)$ which is a contradiction. Thus $G_1 = \pi_1(M_1)$ satisfies condition (i) of Lemma 9.1.

Also, the argument above proves that if $\overline{M_2}$ has hyperbolic interior then for any $g_2 \in \pi_1(\overline{M_2}) \setminus \pi_1(T)$ and $h \in \pi_1(T) \setminus \{1\}$ the condition (ii) is satisfied. So the assertion is proved if $\overline{M_1}$ and $\overline{M_2}$ both have hyperbolic interior with finite volume.

Now consider the case when $\overline{M_2}$ is Seifert fibred. If h is not a power of the element by represented by the normal fiber of $\overline{M_2}$, then it is not in the centralizer of $\pi_1(\overline{M_2})$ and the condition (ii) holds for an appropriate choice of g_2 . Otherwise replace φ with

$$\varphi'(s, t) = \varphi(t, s).$$

Namely, the set of marked points is:

$$\varphi'(s, 1) = \varphi(1, s), 0 \leq s \leq 1$$

and the loop passing through $\varphi'(s, 1)$ is

$$\varphi'_s(t) = \varphi'(s, t) = \varphi(t, s), 0 \leq t \leq 1.$$

Then h , the element of $\pi_1(\overline{M_2})$ represented by loop passing through $\varphi'(1, 1)$, is not in the centralizer of $\pi_1(\overline{M_2})$ and one can choose $g_2 \in \pi_1(\overline{M_2}) \setminus \pi_1(T)$ such that $g_2h \neq hg_2$. This finishes the proof. $\square$

10. PROOF OF MAIN THEOREM: PART I

In this section we prove the first part of the main theorem, namely that if a closed oriented manifold M which is not algebraically hyperbolic, then M or a double cover of M has nontrivial extended loop products.

For this we use the *prime decomposition* and the *theory of characteristic surfaces* for 3-manifolds. We recall some of these theories.

A closed oriented 3-manifold M is said to be *prime* if it cannot be obtained as the connected sum of two 3-manifolds that none of them is homeomorphic to 3-sphere. By the Alexander's theorem S^3 is prime, as every embedded 2-sphere in S^3 bounds a 3-ball.

Prime Decomposition (see [13]): Let M be a compact, connected and oriented 3-manifold. Then there is a decomposition

$$M = P_1 \# P_2 \# \dots \# P_n$$

with each P_i prime and this decomposition is unique up to permutation of P_i 's and insertion or deletion of S^3 's.

A closed 3-manifold is said to be *irreducible* if any embedded 2-sphere S^2 bounds a 3-ball. Again by the Alexander's theorem, S^3 is irreducible. An oriented irreducible 3-manifold is prime. The converse is not true. The only prime oriented 3-manifold which is not irreducible, is $S^1 \times S^2$.

One can deduce a necessary condition for irreducibility from the *sphere theorem*. Here, we state a simplified version of the theorem.

Sphere Theorem (see [6]): Let M be an oriented connected 3-manifold. If $\pi_2(M) \neq 0$ then there is an embedded S^2 in M representing a nontrivial element in $\pi_2(M)$.

The topology of an irreducible 3-manifold M is coarsely determined by the cardinality of the fundamental group. By the *sphere theorem* (see [7]) $\pi_2(M) = 0$. Let $\tilde{M}$ be the universal cover of M . If $\pi_1(M)$ is finite, then $\tilde{M}$ is closed and simply connected. Therefore $\tilde{M}$ is a homotopy sphere. If $\pi_1(M)$ is infinite then $\tilde{M}$ is not closed and we have $\pi_i(\tilde{M}) = 0$ for all i , by Hurewicz theorem. Therefore $\tilde{M}$ is contractible or in other words M is a $K(\pi, 1)$ (aspherical).

Now by the prime decomposition theorem, every 3-manifold M can be written as

$$(10.1) \quad M = (K_1 \# K_2 \# \dots \# K_p) \# (L_1 \# L_2 \# \dots \# L_q) \# (\#_1^r(S^1 \times S^2))$$

where K_i 's are closed aspherical irreducible 3-manifolds with infinite π_1 and L_i are closed 3-manifolds that are finitely covered by a homotopy 3-sphere.

Let S be an embedded surface in a 3-manifold M . We shall say that S is *properly embedded* if $S \cap \partial M = \partial S$. Then S is *two-sided* in M if the normal bundle of S in M is trivial.

A 2 sided surface S in M is said to be *compressible* if either:

- (i) S is a 2-sphere and it bounds a 3-cell in M
- or
- (ii) there is a disk $D \in M$ such that $D \cap S = \partial D$ and ∂D is non contractible in S .

Otherwise S is said to be *incompressible*. An embedded disk is considered to be incompressible. For an incompressible surface S in M , the induced map by inclusion $\pi_1(S) \rightarrow \pi_1(M)$ is injective. Indeed, a 2-sided surface S is incompressible if and only if the map $\pi_1(S) \rightarrow \pi_1(M)$ induced by inclusion is injective¹².

An oriented 3-manifold M , possibly with boundary is called *atoroidal* if any incompressible torus in M is *∂ -parallel*, meaning that it can be isotoped to the boundary of M .

Torus decomposition (see [9] or [10]): For a compact irreducible oriented 3-manifold M there exists a collection $\mathcal{T} \subset M$ of disjoint incompressible tori such that the closure of each component of $M \setminus \mathcal{T}$ is either atoroidal or a Seifert manifold with $\mathcal{T}$ tangent to the fibration. A minimal such a collection $\mathcal{T}$ is unique up to isotopy.

As we are interested in rank 2 abelian subgroups of fundamental groups, torus theorems by Scott ([16]) and Seifert fibred space theorem by Casson-Jungreis [3] and Gabai [6] will be useful to us.

¹² π_1 -injective

Torus Theorem (see [16]): Let M be a compact, connected oriented, irreducible 3-manifold. Suppose that $\pi_1(M)$ has a free abelian subgroup of rank 2. Then either

- (i) M contains a two sided torus such that the inclusion homomorphism $\pi_1(T) \rightarrow \pi_1(M)$ is injective, or
- (ii) $\pi_1(M)$ contains an infinite cyclic normal subgroup.

Seifert Fibred Space Theorem (see [3] or [6]) Let M be a compact connected, oriented, irreducible 3-manifold with infinite π_1 , then M is a Seifert fibred space if and only if $\pi_1(M)$ has a infinite cyclic normal subgroup.

We recall the definition of algebraically hyperbolic manifolds.

Definition An oriented 3-manifold is said to be *algebraically hyperbolic* if it is aspherical ($K(\pi, 1)$) and $\pi_1(M)$ has no rank 2 abelian subgroup.

Proof of the first part of the main theorem: If M is not algebraically hyperbolic then at least one of the following holds:

10.1. If M is not aspherical. By prime decomposition theorem, every 3-manifold M can be written as

$$(10.2) \quad M = (K_1 \# K_2 \# \dots \# K_p) \# (L_1 \# L_2 \# \dots \# L_q) \# (\#_1^r(S^1 \times S^2))$$

where K_i 's are closed aspherical irreducible 3-manifolds with infinite π_1 and L_i are closed 3-manifolds that are finitely covered by a homotopy 3-sphere. If M is not aspherical then one of the following holds:

- (i) If M is a connected sum of manifolds with nontrivial fundamental group then Proposition 6.1 says that M has nontrivial extended loop products.
- (ii) If M has a non separating 2-sphere or in other words there is a $S^2 \times S^1$ factor in its prime decomposition, then by Corollary 3.2 M has nontrivial extended loop products.
- (iii) M has finite fundamental group; Then by Proposition 2.6 M has nontrivial extended loop products.

Thus in either cases M has nontrivial extended loop product.

10.2. If M aspherical and $\mathbb{Z} \oplus \mathbb{Z} \subset \pi_1(M)$. We turn to the case when M is aspherical 3-manifolds with rank 2 abelian groups. It follows from the argument in 10.1 that M is either irreducible or is connected sum of an irreducible 3-manifold and homotopy 3-sphere¹³.

10.2.1. M is irreducible. By Torus theorem and Seifert theorem one of the following holds:

a) M is a closed Seifert manifold. If M is Seifert manifold then by Proposition 4.2 it has nontrivial extended loop products.

b) The collection of tori in the torus decomposition of M is nonempty. This case itself is divided into 3 cases:

1. M contains a non separating torus: If there is a non separating torus in the collection T then by Proposition 3.3 M has nontrivial extended loop products.

¹³Note that such a connected sum is again aspherical.

2. Torus decomposition of M has only Seifert pieces and has no non separating torus: Let $\mathcal{T}$ be the collection of tori provided by the torus decomposition of M and

$$M \setminus \mathcal{T} = M_1 \cup M_2 \dots \cup M_n.$$

$\overline{M_i}$ is a Seifert manifold. Let p_i and b_i be the number of singular fibers and boundary components of $\overline{M_i}$. If S_i , the base surface of $\overline{M_i}$, is orientable then g_i denotes its genus otherwise g_i is number of cross caps in S_i .

If for some i , S_i is orientable and $g_i \geq 1$, then M has a non separating torus and by case 1 we know that M has nontrivial extended loop product. So we assume that $g_i = 0$ if S_i is oriented.

If S_i is orientable then,

$$p_i + b_i \geq 3.$$

To see that suppose that $p_i + b_i \leq 2$. Then one of the following holds:

- (a) $b_i = 1$ and $p_i = 1$;
- (b) $b_i = 2$ and $p_i = 0$

In the first case it follows that $\pi_1(\overline{M_i}) \simeq \mathbb{Z}$ so the boundary components of $\overline{M_i}$ can not be incompressible. In the latter case it follows that $\overline{M_i} \simeq S^1 \times S^1 \times [0, 1]$ this contradicts the minimality of the collection $\mathcal{T}$ as one can extend the Seifert fibration of one of the neighboring $\overline{M_j}$ to $\overline{M_i}$.

Therefore M satisfies all the conditions of Proposition 8.1, hence M or a double cover of M has nontrivial extended loop product.

3. If M has an atoroidal component and has no non separating torus: Suppose that $\mathcal{T}$ is the collection of tori provided by the torus decomposition of M and every torus in $\mathcal{T}$ is separating.

By Thurston theorem ¹⁴[17],

Theorem *A compact, oriented, irreducible atoroidal 3-manifold whose boundary consists of a finite number of tori admits a complete hyperbolic metric of finite volume;*

atoroidal components of $M \setminus \mathcal{T}$ are hyperbolic.

Let M_1 be a hyperbolic component of $M \setminus \mathcal{T}$ and M_2 is another component that is attached to M_1 along a torus T . By proposition 9.2, $M' = \overline{M_1} \cup T_1 \cup \overline{M_2}$ has nontrivial extended loop products.

Consider the same homology classes Γ and H in $\mathbb{H}_*(M')$ constructed in the proof of Proposition 9.2. They can be regarded as elements of $\mathbb{H}_*(M)$. We claim that the loop product

$$p_{A_M}(\Delta(\Gamma)) \bullet p_{A_M}(\Delta H) \neq 0,$$

so M has nontrivial extended loop products.

The calculation of the loop product is the same as of the proof proposition 9.2. Calculating mod 2 only two terms remain, each one an element of $\mathbb{H}_0(M)$ which is represented by a conjugacy class in $\pi_1(M)$. These conjugacy classes are

$$[hg_1g_2] \text{ and } [hg_2g_1].$$

We prove that these two conjugacy classes are different. By the proof of proposition 9.2 we know that hg_1g_2 and hg_2g_1 are not conjugate in $\pi_1(M')$. So by the

¹⁴Geometrization of Haken manifolds.

injectivity lemma (Lemma 7.1), if we show that hg_1g_2 is not conjugate to an element of the fundamental group of one of the boundary components of $\overline{M'}$ it follows that hg_1g_2 and hg_2g_1 are not conjugate in $\pi_1(M)$.

Let $T_1 \cup T_2 \dots \cup T_k = \partial \overline{M'}$ and suppose that hg_1g_2 is conjugate to an element of $\pi_1(T_i)$, $i > 1$ in $\pi_1(M')$. Since one of the

$$T_i \subset \overline{M_1} \text{ or } T_i \subset \overline{M_2}$$

holds let us assume that $T_i \subset \overline{M_1}$. Then

$$\pi_1(T_i) \subset \pi_1(M_1) = \pi_1(\overline{M_1})$$

as T_i is incompressible. If hg_1g_2 is conjugate to an element to $\pi_1(T_i)$ then it is conjugate to an element in $\pi_1(M_1)$. But this is a contradiction since $hg_1g_2 = (hg_1)g_2$ as an element of

$$\pi_1(M') = \pi_1(M_1) *_{\pi_1(T)} \pi_1(M_2)$$

is a cyclically reduced word of length 2 and it is cannot be conjugate to an element of the factor $\pi_1(M_1)$. Now by the injectivity lemma hg_1g_2 and hg_2g_1 are not conjugate in $\pi_1(M)$.

10.2.2. M is a connected sum of an irreducible aspherical manifold and a homotopy 3-sphere. Suppose that $M = N \# P$ where N is irreducible and aspherical and $P \neq S^3$ is a homotopy 3-sphere. M is obtained from N by replacing a standard ball $D \subset N$ with a fake ball D' namely $M = (N \setminus D) \cup_{S^2} D'$. Let

$$f : N \rightarrow M$$

be a homotopy equivalence which sends D to D' and is identity on the complement of D in N . Since $\mathbb{Z} \oplus \mathbb{Z} \subset \pi_1(M)$ hence $\mathbb{Z} \oplus \mathbb{Z} \subset \pi_1(N)$.

By Seifert and Torus theorem either N is Seifert or the collection of tori in its torus decomposition is non empty.

If N has a nontrivial torus decomposition it follows from the previous case, 10.2.1, that N has nontrivial extended loop products. The homology classes in N constructed in 10.2.1 can be regarded as homology classes in M since they can have chain representative which is in the complement of D and the loop products calculated there are also nontrivial in M as $\pi_1(M) \simeq \pi_1(N)$.

If N is a Seifert manifold then one of the 3-homology class constructed in according to Section 4, θ , can not be considered as a homology class in M since it uses all the manifold N but we claim that

$$p_{A_M}(f_{\mathbb{L}}(\theta)) \bullet p_{A_M}(f_{\mathbb{L}}(\alpha)) \neq 0.$$

which proves that M has nontrivial extended loop products.

The proof is exactly the one in Section 4 one just has to notice that we can choose a representative for α such that its marked point is in the complement of D .

11. PROOF OF MAIN THEOREM: PART II, ALGEBRAICALLY HYPERBOLIC MANIFOLDS

In this section we prove the second part of the main theorem, namely that algebraically hyperbolic manifolds have trivial extended loop products.

It follows from Lemma 1.5 that

Lemma 11.1. *If M is a $K(\pi, 1)$ (aspherical) then $\mathbb{L}M$ is homotopy equivalent to*

$$\coprod_{[\alpha] \neq [1] \in \hat{\pi}_1(M)} K(\pi_\alpha, 1)$$

where we choose only one representative for each conjugacy class and π_α is the centralizer of α in $\pi_1(M)$.

Therefore to prove the second part of the theorem we need a good understanding of the centralizers. For that we need following theorem.

Compact Realization Theorem (see [7]): Let M be a connected, oriented 3-manifold, and let K be a subgroup of $\pi_1(M)$ which is finitely generated. Then there is a compact, oriented irreducible 3-manifold M_0 with $\pi_1(M_0) \simeq K$.

Lemma 11.2. *If a closed oriented 3-manifold M is algebraically hyperbolic then the fundamental group of M does not have any nontrivial finite subgroups and the centralizer π_α of an element $\alpha \in \pi_1(M)$ is isomorphic to an additive subgroup of $\mathbb{Q}$.*

Proof. Let $\mathbb{Z}_p$ be a nontrivial finite subgroup of $\pi_1(M)$. Then $K(\mathbb{Z}_p, 1)$ is a covering of M , so it has to be a 3-manifold. But one knows that $K(\mathbb{Z}_p, 1)$ has homology in infinitely many dimensions. So $\pi_1(M)$ has no finite subgroup.

For the second part we first prove that every finitely generated subgroup of π_α is isomorphic to $\mathbb{Z}$. Suppose $\beta_1, \beta_2, \dots, \beta_n$ are elements of π_α . Consider the subgroup $\langle \beta_1, \beta_2, \dots, \beta_n, \alpha \rangle$ which is finitely generated and it is not a free product since it has a nontrivial center. Therefore by compact realization theorem it is isomorphic to the fundamental group of a compact, oriented irreducible 3-manifold. The cyclic subgroup $\langle \alpha \rangle$ is in its center and therefore it is normal. Thus by the Seifert fibred space theorem, the compact realization should be a Seifert manifold. Since $\pi_1(M)$ has no finite subgroup or rank 2 abelian subgroup this Seifert manifold has to be a solid torus¹⁵. As the fundamental group of a solid torus is $\mathbb{Z}$, we have $\langle \beta_1, \beta_2, \dots, \beta_n, \alpha \rangle \simeq \mathbb{Z}$. To prove that π_α is isomorphic to a subgroup of $\mathbb{Q}$, note that since $\pi_1(M)$ is countable and every finitely generated subgroup is cyclic, we can write $\pi_\alpha = G_1 \cup G_2 \dots$ where each G_i is an infinite cyclic subgroup and $G_i \subset G_{i+1}$. Let x_i be a generator for G_i ; then we have $x_i = x_{i+1}^{n_i}$ for some n_i . We construct a map $\varphi : \pi_\alpha \rightarrow \mathbb{Q}$ by letting $\varphi(z_1) = 1$ and $\varphi(z_{i+1}) = \frac{1}{n_1 n_2 \dots n_i}$. This is a well defined map because of the way we chose n_i 's. φ is injective. To see that note that if $x \in \pi_1(M)$ then $x \in G_i$ for some i . So $x = k z_i$ for some i and k . $\varphi(x) = \frac{k}{n_1 n_2 \dots n_i}$. Therefore $\varphi(x) = 0$ implies $k = 0$ or $x = 0$. $\square$

Lemma 11.3. *If G is an additive subgroup of $\mathbb{Q}$ then $K(G, 1)$ has homological dimension one.*

Proof. One can write $G = \varinjlim_{n \rightarrow \infty} G_n$, where G_i 's are cyclic subgroups of G . So $K(G, 1) = \varinjlim_{n \rightarrow \infty} K(G_n, 1)$. Since each $K(G_n, 1) \simeq K(\mathbb{Z}, 1)$ has homological dimension 1 therefore the direct limit $K(G, 1)$ has homological dimension one as taking homology commutes with taking direct limit. $\square$

Proposition 11.4. *If M is algebraically hyperbolic then any finite cover of M has trivial extended loop products.*

¹⁵ M cannot be $S^2 \times S^1$ since it is irreducible.

Proof. Let $\widetilde{M}$ be a finite cover of M . Note that $\widetilde{M}$ is also algebraically hyperbolic. Since $\widetilde{M}$ is a $K(\pi', 1)$ then its free loop space is homotopy equivalent to

$$\coprod_{[\alpha] \in \hat{\pi}'} K(\pi'_\alpha, 1)$$

where we choose only one representative for each conjugacy and π'_α is the centralizer of α in $\pi_1(\widetilde{M})$. So

$$A_{\widetilde{M}} \simeq H_*\left(\coprod_{[\alpha] \in \hat{\pi}', \alpha \neq 1} K(\pi'_\alpha, 1)\right) \simeq \bigoplus_{[\alpha] \in \hat{\pi}', \alpha \neq 1} H_*(K(\pi'_\alpha, 1))$$

Now by Lemma 11.2 and Lemma 11.3, we see that each $K(\pi'_\alpha, 1)$, $\alpha \neq 1$, has homological dimension 1. This proves that the loop product $\bullet$ on $A_{\widetilde{M}}$ is zero because of the absence of homology classes of sufficiently high degree. $\square$

APPENDIX A. SOME FACTS IN FREE PRODUCTS WITH AMALGAMATION

A.1. Some Basic Facts about Free Products with Amalgamation: We recall some definitions and theorems from group theory related to the free product with amalgamation. We follow the terminology and notations of [12] and [11].

Let G_1, G_2 and H be three groups and $\phi_i : H \rightarrow G_i$, $i = 1, 2$, be two injective homomorphisms. We can regard H as a subgroup of each G_i , $i = 1, 2$ and we identify each element of H with its image under ϕ_i while we are in G_i . So if $h \in H$, then h can be considered as an element of both G_1 and G_2 . Following [11], we can form the amalgamated free product $G_1 *_H G_2$ of G_1 and G_2 over H .

A sequence of elements $g_1, g_2, \dots, g_n \in G_1 *_H G_2$ is called *reduced* if:

- (1) Each g_i is in one of the factors G_1 or G_2 ;
- (2) g_i, g_{i+1} come from different factors; So if $n > 1$ then $g_i \notin H$ for all i .
- (3) If $n = 1$ then $g_1 \neq 1$.

$g_1 g_2 \dots g_n \in G_1 *_H G_2$ is called a *reduced word* if $g_1, g_2, \dots, g_n$ is a reduced sequence.

There is no canonical way of presenting the elements of $G_1 *_H G_2$ uniquely. Once a choice of left coset representatives for G_1/H and G_2/H is made then every element can be represented uniquely. Let $c_1, \dots, c_n$ be a right coset representative for G_1/H and G_2/H . Here is an informal description of the representation:

Let g be an element in $G_1 *_H G_2$ and $g = g_1 g_2 \dots g_r$ be a reduced presentation of g . Start with g_r , there is an $h_r \in H$ and $c_r \in G_1/H$ or G_2/H (depending in which factor is g_r) such that

$$g_r = h_r c_r.$$

Now by replacing g_r by $h_r c_r$ we have $g = g_1 g_2 \dots g_{r-1} h_r c_r$. Again there is a $h_{r-1} \in H$ and c_{r-1} such that

$$g_{r-1} h_r = h_{r-1} c_{r-1}.$$

By replacing $g_{r-1} h_r$ by $h_{r-1} c_{r-1}$ we have $g = g_1 g_2 \dots g_{r-2} h_{r-1} c_{r-1} c_r$. Continuing this procedure we end up with $g = h_1 c_1 c_2 \dots c_r$. This is the desired presentation. While stating it more precisely, the following theorem also asserts its uniqueness (*i.e.* its independence from the reduced word we used in the construction).

Theorem A.1. (See [12] pp.201) Let $G = G_1 *_H G_2$ where H is a mutual subgroup of G_1 and G_2 . Suppose a specific right coset representative systems for G_1/H and G_2/H have been selected. Then with each element of g of G we can associate a unique sequence¹⁶

$$(h, c_1, c_2, \dots, c_r)$$

such that:

- (i) h is an element, possibly 1, of H ;
- (ii) c_i is a coset representative of G_1/H or G_2/H ;
- (iii) $c_i \neq 1$
- (iv) c_i and c_{i+1} are not both in G_1 or G_2 .
- (v) If h' and c'_i s are the images of h or c_i 's under the homomorphisms of G_1 or G_2 into G , then $g = h'c'_1c'_2\dots c'_r$ in G .

A sequence of elements $g_1, g_2, \dots, g_n \in G_1 *_H G_2$ is called *cyclically reduced* if all cyclic permutations of the sequence $g_1, g_2, \dots, g_n$ are also reduced.

$g_1g_2\dots g_n \in G_1 *_H G_2$ is called a *cyclically reduced word* if $g_1, g_2, \dots, g_n$ is a cyclically reduced sequence.

Also every element of $G_1 *_H G_2$ is conjugate to $g_1g_2\dots g_n$ where the sequence $g_1, g_2, \dots, g_n$ is cyclically reduced. Indeed if $g = g'_1g'_2\dots g'_k$ where $g'_1, g'_2, \dots, g'_k$ is a reduced sequence. If g'_1 and g'_k are from different factors, then clearly $g'_1g'_2\dots g'_k$ is cyclically reduced. Otherwise consider $g'_kg'_k{}^{-1} = g'_kg'_1g'_2\dots g'_{k-1}$ and replace $g'_kg'_1$ by $g' = g'_kg'_1$ which is from the same factor. Now $g', g'_2, \dots, g'_{k-1}$ is cyclically reduced.

Theorem A.2. (see [11] p.187) If $c_1, c_2, \dots, c_n$, $n \geq 1$, is a reduced sequence in $G = G_1 *_H G_2$, then the product $c_1c_2\dots c_n \neq 1$ in G .

The following theorem describes the conjugacy classes of $G_1 *_H G_2$.

Theorem A.3. (see [12] pp.212)

Let $G = G_1 *_H G_2$. Then every element of G is conjugate to a cyclically reduced element of G . Moreover, suppose that g is a cyclically reduced element of G . Then:

- (i) If g is conjugate to a cyclically reduced word $p_1p_2\dots p_r$ where $r \geq 2$; then g can be obtained by cyclically permuting $p_1, p_2, \dots, p_r$ and then conjugating by an element of H .
- (ii) If g is conjugate to an element h in H , then g is in some factor and there is a sequence of $h, h_1, \dots, h_t, g$ where h_i is in H and consecutive terms of the sequence are conjugate in some factor.
- (iii) If g is conjugate to an element g' in some factor but not in a conjugate of H then g and g' are in the same factor and are conjugate in that factor.

Lemma A.4. Suppose that G_1, G_2 and H are three groups with monomorphisms $\phi_i : H \rightarrow G_i$, $i = 1, 2$. Let $G_1 *_H G_2$ denote the amalgamated free product of G_1 and G_2 over H . Let $g_1 \in G_1 \setminus H$ and $g_2 \in G_2 \setminus H$ and $h \neq 1$ in H such that:

- (i) $g_1^{-1}Hg_1 \cap H = 1$

¹⁶This presentation is called *reduced form* (see [12]).

(ii) $g_2h \neq hg_2$.

Then two conjugacy classes $[hg_1g_2]$ and $[hg_2g_1]$ of $G_1 *_H G_2$ are different.

Proof. Suppose that $[hg_1g_2] = [hg_2g_1]$. It is clear that

$$(A.1) \quad [hg_1g_2] = [g_2hg_1]$$

$$(A.2) \quad [hg_2g_1] = [g_2g_1h]$$

So $g_2(hg_1)$ is conjugate to $g_2(g_1h)$. Since $g_2(g_1h)$ is cyclically reduced (of length 2, $p_1 = g_2$ & $p_2 = g_1h$) by theorem A.3 part (ii), $g_2(g_1h)$ is conjugate to $g_2(g_1h)$ or $(g_1h)g_2$ by an element in H .

In the latter case we have

$$g_2(hg_1) = h_1(g_1h)g_2h_1^{-1}$$

which is equivalent to

$$g_2^{-1}(h_1g_1h)^{-1}g_2(hg_1h_1) = 1.$$

But this contradicts Theorem A.2. Since the sequence

$$g_2^{-1}, (h_1g_1h)^{-1}, g_2, (hg_1h_1)$$

is a reduced sequence.

Therefore there should be an element $h_1 \in H$ such that

$$g_2hg_1 = h_1g_2g_1hh_1^{-1}$$

or equivalently

$$g_2hg_1h_1 = h_1g_2g_1h$$

or

$$(A.3) \quad (g_2h)(g_1h_1) = (h_1g_2)(g_1h)$$

Now using the uniqueness of representation in terms of left cosets of H Theorem 1 (also see the construction) the equation (A.3) implies that g_1h_1 and g_1h are in the same left coset in G_1/H or in other words, there exists $h_2 \in H$ such that

$$g_1h_1 = h_2g_1h$$

or equivalently

$$h_1h^{-1} = g_1^{-1}h_2g_1$$

So $h_1h^{-1} = g_1^{-1}h_2g_1 \in g_1^{-1}Hg_1 \cap H = 1$ hence $h_2 = h_1h^{-1} = 1$ which implies $h_1 = h$. By substituting $h_1 = h$ in (A.3) and simplifying, it follows

$$g_2h = hg_2$$

which is a contradiction. Therefore $[hg_1g_2] \neq [hg_2g_1]$

□

A.2. Tree of Groups. A *tree of groups* (G, T) consists of a finite tree T and G a collection of groups

$$G = \{G_e\}_{e \in \text{edge}T} \coprod \{G_v\}_{v \in \text{vert}T},$$

a group G_v for every vertex $v \in \text{vert}T$, a group G_e for every edge e in $\text{edge}T$ and a monomorphism

$$f_e^v : G_e \rightarrow G_v$$

if v is a vertex of the edge e and

We call G_v a *vertex group* if v is a vertex of T and G_e a *edge group* if e is an edge of T . If v is a vertex of the edge e then G_e can be considered as a subgroup of G_v .

Suppose that (G, T) is a tree of groups, v_0 a vertex of T . Let $n = |\text{edge}T|$. Consider a sequence of trees T_i , $1 \leq i \leq n$ such that

- (i) $|\text{edge}T_i| = i$, $0 \leq i \leq n$
- (ii) $\text{vert}T_0 = \{v_0\}$,
- (iii) $T_i \subset T_{i+1}$ and
- (iv) $T_n = T$.

Let $e_1, e_2, \dots, e_n$ be the sequences of the edges and $v_1, v_2, \dots, v_n$ be the sequence of vertices such that

- $\text{edge}T_1 = \{e_1\}$
- $\text{vert}T_1 = \{v_0, v_1\}$
- $\text{edge}T_{i+1} = \text{edge}T_i \cup \{e_{i+1}\}$, $1 \leq i \leq n-1$.
- $\text{vert}T_{i+1} = \text{vert}T_i \cup \{v_{i+1}\}$ $1 \leq i \leq n-1$.

Let

$$\begin{aligned} G_{T_0} &= G_{v_0}, \\ &\text{and} \\ G_{T_i} &= G_{T_{i-1}} *_{G_{e_n}} G_{v_n} \\ &1 \leq i \leq n \end{aligned}$$

$G_T = G_{T_n}$ is called the amalgamation of G_v 's along G_e 's and is independent of the choice of the sequence T_i and depends only on (G, T) and there is an inclusion

$$\varphi_k : G_k \rightarrow G_T$$

where k is a vertex or an edge.

The following lemma is a generalization of theorem A.3 part (iii).

Lemma A.5. *Let (G, T) be a tree of groups and v_0 a vertex of T . Consider the construction above. Suppose that $[a]$ and $[b]$ are two distinct conjugacy classes of G_{v_0} and a is not conjugate in G_{v_0} to an element of any edge group $G_e \subseteq G_{v_0}$ where v_0 is a vertex of e . Then $[a]$ and $[b]$ as the conjugacy classes of G_T are distinct.*

Proof. T_i 's, v_i 's and e_i 's are as above.

We prove the lemma by induction. We prove that for every i , $1 \leq i \leq n$

- (1) $[a]$ and $[b]$ are distinct as conjugacy classes of G_{T_i} .
- (2) a is not conjugate in G_{T_i} to an element in the vertex group G_w , where $w \neq v_0$ is a vertex of T_i .

We verify the statement for $i = 1$. By the assumption and part (iii) of theorem A.3 it follows that the first statement of induction is true for $i = 1$. For the second part, suppose that a is conjugate in G_{T_1} to an element of G_{v_1} , then again by the third part of theorem A.3, a has to be conjugate to an element of G_{e_1} and then by part (ii) it has to be conjugate in G_{v_0} to an element of the vertex group G_{e_1} which contradicts our assumption.

Suppose that the statement is true for i . By the second statement of the induction for i , a is not conjugate in G_{T_i} to an element of $G_{e_{i+1}} \subseteq G_{T_i}$. Therefore by part (iii), a and b represent different conjugacy classes in $G_{T_{i+1}}$ as

$$G_{T_{i+1}} = G_{T_i} *_{G_{e_{i+1}}} G_{v_{i+1}},$$

proving the first statement of the induction for $i+1$. To prove the second statement for $i+1$, suppose that by contrary a is conjugate to an element of G_w , where $w \neq v_0$ and $w \in T_{i+1}$.

Since

$$G_{T_{i+1}} = G_{T_i} *_{G_{e_{i+1}}} G_{v_{i+1}},$$

and a is not conjugate in G_{T_i} to an element of $G_{e_{i+1}}$ then again by part (iii) of theorem 3 we must have $w \in T_i$ and a is conjugate in G_{T_i} to an element of G_w . This contradicts the assumption of the induction for i . Hence the second statement of the induction holds for $i+1$.

Finally for $i = n$ we get the statement of the lemma. $\square$

APPENDIX B. FINITE COVERS OF CERTAIN SEIFERT MANIFOLDS WITH BOUNDARY

This appendix is devoted to explaining some facts that are used in the proof of main theorem, section 9. We follow the notations introduced section 8.

Construction 1: Suppose M is a compact orientable Seifert manifold with $g = 0$, $p = 1$ and $b = 2$ and where the singular fiber has type $(2, 1)$.

Then M has a double cover M' with $g' = 0$, $p' = 0$ and $b' = 3$ boundary components and two of the boundary components are identified under the covering map :





FIGURE 8. N

Consider $D \subset \mathbb{R}^2$ the unit disk centered at the origin and F a union of two open disks of radius $1/3$ centered at $(1/2, 0)$ and $(-1/2, 0)$ (Figure 8).

Let $M' = N \times \mathbb{S}^1$ where

$$N = D \setminus F.$$

$\mathbb{Z}_2$ acts on M' freely where the action on both factors is realized by 180° rotation.

FIGURE 9. Double cover M' with 3 boundary components


Indeed,

$$M = M' / \mathbb{Z}^2$$

and M' is the desired double covering (Figure 9). M' has 3 boundary components, of which two are identified under the covering map. $\square$

Construction 2: Let M be a compact orientable Seifert manifold with $g = 0$, $b = 1$ and $p = 2$ singular fibers of type $(2, 1)$ and (r, s) . Then M has a double cover M' with $g' = 0$, $b' = 1$ and $p' = 2$ and the singular fibers are of type (r, s) :

Let N be the 2-sphere $S^2 \setminus D \subset \mathbb{R}^3$, where D is a set of n open disks located on the equator in xy -plane which is invariant under $2\pi/s$ rotation around z -axis and 180° rotation; and If s is even then D does not meet the x axis otherwise D meets the x -axis exactly at one point.

FIGURE 10. Double cover $\widetilde{M}$

The $2s$ -Dihedral group,

$$D_{2s} = \langle a, b \mid a^2 = 1, b^s = 1, aba = b^{-1} \rangle$$

acts on $M' = N \times \mathbb{S}^1$ freely (see Figure 10);

- (i) a acts by 180° rotation about x -axis in the first factor and 180° rotation in the second factor.
- (ii) b acts by $2\pi/s$ rotation in the first factor and $2r\pi/s$ rotation in the second factor.

Let $M' = N / \langle b \rangle$, then M' is a Seifert manifold with two singular fibers of type (r, s) and one boundary component and base surface a disk. $\langle a \rangle$ acts on M' freely since $\langle b \rangle$ is normal subgroup of D_{2s} . Indeed,

$$M = M' / \langle a \rangle$$

Construction 3: Let M be a compact Seifert manifold with $g = 0$, $b = 1$ and $p = 2$ singular fibers of type $(2, 1)$. Then M has a double cover M' with $g' = 0$, $b' = 2$ and $p' = 0$ and the boundary components of M are identified homeomorphically under the covering map:

Consider the annulus $C = S^1 \times [-1, 1] \subset \mathbb{R}^3$, where S^1 is the unit circle in (x, y) plane and centered at the origin (Figure 11).




FIGURE 11. Annulus $C \subset \mathbb{R}^3$

$\mathbb{Z}_2$ acts on C with a and b as the only fixed points. The action is realized by 180° rotation about y axis.

$\mathbb{Z}_2$ acts on

$$M' = C \times \mathbb{S}^1$$

freely where the action on the first factor as described above and on the second factor is rotation by 180° . Indeed,

$$M \simeq M' / \mathbb{Z}_2$$

and the boundary components of M' are identified under homeomorphically.


Lemma B.1. *Let M be a closed oriented 3-manifold and T be the collection of tori provided by torus decomposition of M . Suppose that $M \setminus T = M_1 \cup M_2 \cup \dots \cup M_n$ with all M_i 's Seifert manifold such that:*

- (1) $n \geq 3$ and $g_i = 0$ for all i 's;
- (2) $b_1 = b_n = 1$ and $b_i = 2$ for $i \neq 1, n$;
- (3) $p_1 = 2$, $A_1 = \{2, r\}$;
- (4) $p_2 = 1$ and $A_2 = \{2\}$;

Then M has a double cover M whose torus decomposition has a Seifert component with 3 boundary components.

Proof. Let M'_1 be the double cover of $\overline{M_1}$ provided by Construction 2 and M'_2 be the double cover of M_2 by Construction 1. M'_1 has one boundary S component with an induced involution by the covering map. M'_2 has 3 boundary components T_1 , T_2 and T_3 where T_2 and T_3 are identified under the covering map and T_1 has an induced involution.

Glue M'_1 and M'_2 along S and T_1 and then two copies of $\overline{M_3} \cup \overline{M_4} \dots \overline{M_n}$ to $M'_1 \cup_{S=T_1} M'_2$ along T_2 and T_3 in such a way that the result is a covering of M . This is the double covering which has a Seifert component with two singular fibers of multiplicity r (Figure 12). $\square$

FIGURE 12. Double cover $\widetilde{M}$

Lemma B.2. *Let M be a closed oriented 3-manifold and T be the collection of tori provided by torus decomposition of M . Suppose that $M \setminus T = M_1 \cup M_2$ with all M_i 's Seifert manifold such that:*

- (1) $b_1 = 1$, $p_1 = 2$ and $A_1 = \{2, r\}$;
- (2) $b_2 = 1$, $p_2 = 2$ and $A_2 = \{2, s\}$.

Then M has a double cover whose torus decomposition 2 Seifert components with 2 singular fibers of multiplicity r and s .

Proof. Let M'_1 and M'_2 be 2-covers of $\overline{M}_1$ and $\overline{M}_2$ provided by Construction 2. M'_1 and M'_2 have one boundary component where the covering map induces an involution on the boundary. Glue M'_1 and M'_2 along the boundary in such a way that the result is a double cover of M . This is the desired 2-cover. $\square$

Lemma B.3. *Let M be a closed oriented 3-manifold and T be the collection of tori provided by torus decomposition of M . Suppose that $M \setminus T = M_1 \cup M_2 \cup \dots \cup M_n$ with all M_i 's Seifert manifold; and*


- (1) $b_1 = b_n = 1$; $b_i = 2$ for $i \neq 1, n$;
- (2) $p_1 = p_n = 2$;
- (3) $A_1 = A_n = \{2\}$.

Then M has a double cover with a non separating torus.

Proof. The desired 2-cover of M is constructed as follows: Let $M'_1 \xrightarrow{f_1} \overline{M}_1$ and $M'_n \xrightarrow{f_2} \overline{M}_n$ be the double covers of $\overline{M}_1$ and $\overline{M}_n$ provided by Construction 3. Consider two copies of $M \setminus (M_1 \cup M_n)$ and glued them to $M'_1 \cup M'_n$ so that the result $\widetilde{M}$ double-covers M (Figure 13) in such a ways that covering map restricted to M'_1 and M'_n is f_1 and f_2 and on the rest of $\widetilde{M}$ it is the trivial covering map. This can be done since the covering map f_1 (f_n) identifies different components of M'_1 (resp. M'_n). $\square$

REFERENCES

- [1] Martin Bendersky , Moira Chas & Dennis Sullivan , *Private Communication*, CUNY Graduate Center.
- [2] Benedetti R. & Petronio C., *Lectures on Hyperbolic Geometry*, Universitext Springer-Verlag, Berlin, **1991**.

FIGURE 13. Double cover $\widetilde{M}$ with a non separating torus

- [3] A. Casson, D. Jungreis, *Convergence groups and Siefert fibered 3-manifolds*, Invent. Math. **118** (1994) 441-456.
- [4] M. Chas, *Combinatorial Lie bialgebras of curves on surfaces*, Preprint archive xxx.lanl.gov **GT/0105178**, to appear in Topology.
- [5] M. Chas, D. Sullivan, *String Topology*, Preprint archive, xxx.lanl.gov, **GT/9911159**
- [6] D. Gabai, *Convergence groups are Fuchsian groups*, Ann. of math.(2) **92** 136 (1992) 447-510.
- [7] J. Hempel, *3-manifolds*, Annals of Mathematics studies, no. 48, Princeton University Press, N.J., 1976.
- [8] W. Jaco, *Lectures on Three-Manifold Topology*, CBMS Regional Conference Series in Mathematics, 43. American Mathematical Society, Providence, R.I., 1980.
- [9] W. Jaco, P. Shalen, *Seifert fibered spaces in 3-manifolds*, Mem. Amer. Math. Soc. **21** (1979), no. 220.
- [10] K. Johannson, *Homotopy equivalence of 3-manifolds with boundaries*, Lecture Notes in Mathematics, 761. Springer, Berlin, 1979 ISBN: 3-540-09714-7 (Reviewer: John Hempel), Springer, Berlin.
- [11] R. Lyndon, P. Schupp, *Combinatorial Group Theory*, Reprint of the 1977 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001. ISBN: 3-540-41158-5
- [12] W. Magnus, A. Karrass, D. Solitar, *Combinatorial group Theory*, 1966, John Wiley & Sons, Inc.
- [13] J. Milnor, *A unique factorization for 3-manifolds*, Amer. J. Math. **84** (1962), 1-7
- [14] J. P. Serre, *Trees*, Trees. Translated from the French by John Stillwell. Springer-Verlag, Berlin-New York, ISBN: 3-540-10103-9 1980
- [15] G.P. Scott, *Finitely generated 3-manifold groups are finitely presented.*, J. London Math. Soc. (2) **6**(1973) 437-440.
- [16] P. Scott, *A new proof of annulus and torus theorem*, Amer. J. Math., **102** (1980), no. 2, 241-277.
- [17] Thurston W., *Three dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S)) **6**, (1982), no. 3, 357-381.

CENTRE DE MATHÉMATIQUE, ECOLE POLYTECHNIQUE, PALAISEAU 91128, FRANCE.

E-mail address: habbaspour@math.polytechnique.fr